

BEST AVAILABLE COPY

PCT/EP200 4 / 0 5 1 6 2 5



**Europäisches
Patentamt**

**European
Patent Office**

**Office européen
des brevets**

REC'D 04 OCT 2004

WIPO

PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03102313.8

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1 (a) OR (b)

Der Präsident des Europäischen Patentamts;
im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk



PCT/EP2004/051829

Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

Anmeldung Nr.:
Application no.: 03102313.8
Demande no:

Anmeldetag:
Date of filing: 28.07.03
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Applied Research Systems ARS Holding N.V.
Pietermaai 15
Curacao
ANTILLES NEERLANDAISES

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Imino-azolinone-vinyl fused-benzene derivatives

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

G07D233/00

An Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of
filing/Etats contractants désignés lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT RO SE SI SK TR LI

Imino-azolinone-vinyl fused-benzene derivatives

Field of the invention

This present invention is related to the use of imino-azolinone-vinyl fused-benzene derivatives of formula (I) for the treatment and/or prophylaxis of autoimmune disorders and/or inflammatory diseases, cardiovascular diseases, neurodegenerative diseases, bacterial or viral infections, allergy, asthma, pancreatitis, multiorgane failure, kidney diseases, platelet aggregation, cancer, sperm motility, graft rejection or lung injuries. Specifically, the present invention is related to substituted imino-azolinone-vinyl fused-benzene derivatives for the modulation, notably the inhibition of the activity or function of the phospho-inositide-3'OH kinase family, PI3K, particularly of PI3K γ .

Background of the invention

Cellular plasma membranes can be viewed as a large store of second messenger that can be enlisted in a variety of signal transduction pathways. As regards function and regulation of effector enzymes in phospholipid signalling pathways, these enzymes generate second messengers from the membrane phospholipid pool (class I PI3 kinases (e.g. PI3K γ)) are dual-specific kinase enzymes, means they display both: lipid kinase (phosphorylation of phospho-inositides) as well as protein kinase activity, shown to be capable of phosphorylation of other protein as substrates, including auto-phosphorylation as intra-molecular regulatory mechanism. These enzymes of phospholipid signalling are activated in response to a variety of extra-cellular signals such as growth factors, mitogens, integrins (cell-cell interactions) hormones, cytokines, viruses and neurotransmitters such as described in Scheme 1 hereinafter and also by intra-cellular cross regulation by other signaling molecules (cross-talk, where the original signal can activate some parallel pathways that in a second step transmitt signals to PI3Ks by intra-cellular signaling events), such as small GTPases, kinases or phosphatases for example.


The inositol phospholipids (phosphoinositides) intracellular signalling pathway begins with binding of a signalling molecule (extracellular ligands, stimuli, receptor dimerization, transactivation by heterologous receptor (e.g. receptor tyrosine kinase)) to a G-protein linked transmembrane receptor integrated into the plasma membrane.

- 5 PI3K converts the membrane phospholipid PIP(4,5)2 into PIP(3,4,5)3 which in turn can be further converted into another 3' phosphorylated form of phosphoinositides by 5'-specific phospho-inositide phosphatases, thus PI3K enzymatic activity results either directly or indirectly in the generation of two 3'-phosphoinositide subtypes that function as 2nd messengers in intra-cellular signal transduction (*Trends Biochem Sci.* 22(7) p.267-72
10 (1997) by Vanhaesebroeck B et al., *Chem Rev.* 101(8) p.2365-80 (2001) by Leslie N.R et al (2001); *Annu Rev Cell Dev Biol.* 17 p.615-75 (2001) by Katso R. et al. and *Cell Mol Life Sci.* 59(5) p.761-79 (2002) by Tokerau et al.). Multiple PI3K isoforms categorized by their catalytic subunits, their regulation by corresponding regulatory subunits, expression patterns and signaling-specific functions (p110 α , β , δ , and γ) perform this enzymatic
15 reaction (*Exp Cell Res.* 25(1) p.239-54 (1999) by Vanhaesebroeck B. and *Annu Rev Cell Dev Biol.* 17 p.615-75 (2001) by Katso R. et al).

- The evolutionary conserved isoforms p110 α and β are ubiquitously expressed, while δ and γ are more specifically expressed in the haematopoietic cell system, smooth muscle cells, myocytes and endothelial cells (*Trends Biochem Sci.* 22(7) p.267-72 (1997) by
20 Vanhaesebroeck B et al.). Their expression might also be regulated in an inducible manner depending on the cellular-, tissue type and stimuli as well as disease context.

- To date, eight mammalian PI3Ks have been identified, divided into three main classes (I, II, and III) on the basis of sequence homology, structure, binding partners, mode of activation, and substrate preference *in vitro*. Class I PI3Ks can phosphorylate phosphatidylinositol
25 (PI), phosphatidylinositol-4-phosphate, and phosphatidylinositol-4,5-bisphosphate (PIP2) to produce phosphatidylinositol-3-phosphate (PIP), phosphatidylinositol-3,4-bisphosphate, and phosphatidylinositol-3,4,5-trisphosphate, respectively. Class II PI3Ks phosphorylate PI and

phosphatidylinositol-4-phosphate. Class III PI3Ks can only phosphorylate PI (*Trends Biochem Sci.* **22**(7) p.267-72 (1997) by Vanhaesebroeck B et al, *Exp Cell Res.* **25**(1) p.239-54 (1999) by Vanhaesebroeck B. and *Chem Rev.* **101**(8) p.2365-80 (2001) by Leslie N.R et al (2001)) G-protein coupled receptors mediated phosphoinositide 3'OH-kinase activation
 5 via small GTPases such as G $\beta\gamma$ and Ras, and consequently PI3K signaling plays a central role in establishing and coordinating cell polarity and dynamic organization of the cytoskeleton - which together provides the driving force of cells to move.



Scheme A

10 As above illustrated in Scheme 1, Phosphoinositide 3-kinase (PI3K) is involved in the phosphorylation of Phosphatidylinositol (PtdIns) on the third carbon of the inositol ring.

The phosphorylation of PtdIns to 3,4,5-triphosphate (PtdIns(3,4,5)P₃), PtdIns(3,4)P₂ and PtdIns(3)P act as second messengers for a variety of signal transduction pathways, including those essential to cell proliferation, cell differentiation, cell growth, cell size, cell survival, apoptosis, adhesion, cell motility, cell migration, chemotaxis, invasion,

5 cytoskeletal rearrangement, cell shape changes, vesicle trafficking and metabolic pathway (*Annu Rev Cell Dev Biol.* 17 p.615-75 (2001) by Katso et al. and *Mol Med Today* 6(9) p.347-57 (2000) by Stein R.C). Chemotaxis – the directed movement of cells toward a concentration gradient of chemical attractants, also called chemokines is involved in many important diseases such as inflammation/auto-immunity, neurodegeneration, angiogenesis,

10 invasion/metastasis and wound healing (*Immunol Today* 21(6) p.260-4 (2000) by Wyman NP et al.; *Science* 287(5455) p.1049-53 (2000) by Hirsch et al.; *FASEB J* 15(11) p.2019-21 (2001) by Hirsch et al. and *Nat Immunol.* 2(2) p.108-15 (2001) by Gerard C. et al.).

Recent advances using genetic approaches and pharmacological tools have provided insights into signaling and molecular pathways that mediate chemotaxis in response to

15 chemoattractant activated G-protein coupled receptors PI3-Kinase, responsible for generating these phosphorylated signalling products, was originally identified as an activity associated with viral oncoproteins and growth factor receptor tyrosine kinases that phosphorylates phosphatidylinositol (PI) and its phosphorylated derivatives at the 3'-hydroxyl of the inositol ring (Panayotou et al., *Trends Cell Biol.* 2 p.358-60 (1992)).

20 However, more recent biochemical studies revealed that, class I PI3 kinases (e.g. class IB isoform PI3K γ) are dual-specific kinase enzymes, means they display both: lipid kinase (phosphorylation of phospho-inositides) as well as protein kinase activity, shown to be capable of phosphorylation of other protein as substrates, including auto-phosphorylation as intra-molecular regulatory mechanism.

25 PI3-kinase activation, is therefore believed to be involved in a range of cellular responses including cell growth, differentiation, and apoptosis (Parker et al., *Current Biology*, 5 p.577-99 (1995), Yao et al., *Science*, 267 p.2003-05 (1995)). PI3-kinase appears to be

involved in a number of aspects of leukocyte activation. A p85-associated PI3-kinase activity has been shown to physically associate with the cytoplasmic domain of CD28, which is an important costimulatory molecule for the activation of T-cells in response to antigen (Pages et al., *Nature*, 369 p.327-29 (1994); Rudd, *Immunity* 4 p.527-34 (1996)).

5 Activation of T cells through CD28 lowers the threshold for activation by antigen and increases the magnitude and duration of the proliferative response. These effects are linked to increases in the transcription of a number of genes including interleukin-2 (IL2), an important T cell growth factor (Fraser et al., *Science*, 251 p.313-16 (1991)). Mutation of CD28 such that it can no longer interact with PI3-kinase leads to a failure to initiate IL2

10 production, suggesting a critical role for PI3-kinase in T cell activation. PI3K γ has been identified as a mediator of G beta-gamma-dependent regulation of JNK activity, and G beta-gamma are subunits of heterotrimeric G proteins (*J. Biol. Chem.* 273(5) p.2505-8 (1998)). Cellular processes in which PI3Ks play an essential role include suppression of apoptosis, reorganization of the actin skeleton, cardiac myocyte growth, glycogen synthase

15 stimulation by insulin, TNF α -mediated neutrophil priming and superoxide generation, and leukocyte migration and adhesion to endothelial cells.


Recently, (*Immunity* 16(3) p.441-51 (2002)) it has been described that PI3K γ relays inflammatory signals through various G(i)-coupled receptors and its central to mast cell function, stimuli in context of leukocytes, immunology includes cytokines, chemokines,

20 adenosines, antibodies, integrins, aggregation factors, growth factors, viruses or hormones for example (*J. Cell. Sci.* 114(Pt 16) p.2903-10 (2001) by Lawlor MA et al., *Immunity* 16(3) p.441-51 (2002) by Laffargue M. et al. and *Curr. Opin. Cell Biol.* 14(2) p.203-13 (2002) by Stephens L. et al.).


Specific inhibitors against individual members of a family of enzymes provide invaluable

25 tools for deciphering functions of each enzyme. Two compounds, LY294002 and wortmannin (cf. hereinafter), have been widely used as PI3-kinase inhibitors. These compounds are non-specific PI3K inhibitors, as they do not distinguish among the four

members of Class I PI3-kinases. For example, the IC_{50} values of wortmannin against each of the various Class I PI3-kinases are in the range of 1-10 nM. Similarly, the IC_{50} values for LY294002 against each of these PI3-kinases is about 15-20 μ M (Fruman et al., *An. Rev. Biochem.*, 67 p.481-507 (1998)), also 5-10 microM on CK2 protein kinase and some inhibitory activity on phospholipases. Wortmannin is a fungal metabolite which irreversibly inhibits PI3K activity by binding covalently to the catalytic domain of this enzyme. Inhibition of PI3K activity by wortmannin eliminates the subsequent cellular response to the extracellular factor. For example, neutrophils respond to the chemokine fMet-Leu-Phe (fMLP) by stimulating PI3K and synthesizing $PtdIns(3, 4, 5)P_3$. This synthesis correlates with activation of the respirators burst involved in neutrophil destruction of invading microorganisms. Treatment of neutrophils with wortmannin prevents the fMLP-induced respiratory burst response (Thelen et al. *PNAS* 91 p.4960-64 (1994)). Indeed, these experiments with wortmannin, as well as other experimental evidence, shows that PI3K activity in cells of hematopoietic lineage, particularly neutrophils, monocytes, and other types of leukocytes, is involved in many of the non-memory immune response associated with acute and chronic inflammation.



LY 294002



Wortmannin


Based on studies using wortmannin, there is evidence that PI3-kinase function is also required for some aspects of leukocyte signaling through G-protein coupled receptors (Thelen et al., *Proc. Natl. Acad. Sci. USA*, 91 p.4960-64 (1994)). Moreover, it has been shown that wortmannin and LY294002 block neutrophil migration and superoxide release. However, in as much as these compounds do not distinguish among the various isoforms of

PI3K, it remains unclear which particular PI3K isoform or isoforms are involved in these phenomena.

Cyclooxygenase inhibiting benzofuran derivatives are disclosed by John M. Janusz et al., in *J.Med.Chem.* 1998; Vol 41, No. 18.

5 Summary of the invention

The present invention relates to the use of 2-imino-azolinone-vinyl fused-benzene derivatives of formula (I):



- A, X, Y, R¹, R² and G of formula (I) are defined in the below detailed description. The compounds of formula (I) are useful as medicaments in particular for the treatment and/or prophylaxis of autoimmune disorders and/or inflammatory diseases, cardiovascular diseases, neurodegenerative diseases, bacterial or viral infections, kidney diseases, platelet aggregation, cancer, transplantation, graft rejection or lung injuries. According to one embodiment of the present invention, the compounds of formula (I) are inhibitors of phosphato-inositides 3-kinases (PI3Ks), particularly of Phosphatoinositides 3-kinases gamma (PI3K γ).

Detailed description of the invention:

The following paragraphs provide definitions of the various chemical moieties that make up the compounds according to the invention and are intended to apply uniformly through-

out the specification and claims unless an otherwise expressly set out definition provides a broader definition.

5 "C₁-C₆-alkyl" refers to monovalent alkyl groups having 1 to 6 carbon atoms. This term is exemplified by groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, *tert*-butyl, n-hexyl and the like.

"Aryl" refers to an unsaturated aromatic carbocyclic group of from 6 to 14 carbon atoms having a single ring (e.g., phenyl) or multiple condensed rings (e.g., naphthyl). Preferred aryl include phenyl, naphthyl, phenantrenyl and the like.

10 "C₁-C₆-alkyl aryl" refers to C₁-C₆-alkyl groups having an aryl substituent, including benzyl, phenethyl and the like.

"Heteroaryl" refers to a monocyclic heteroaromatic, or a bicyclic or a tricyclic fused-ring heteroaromatic group. Particular examples of heteroaromatic groups include optionally substituted pyridyl, pyrrolyl, furyl, thienyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,3,4-triazinyl, 1,2,3-triazinyl, benzofuryl, [2,3-dihydro]benzofuryl, isobenzofuryl, benzothienyl, benzotriazolyl, isobenzothienyl, indolyl, isoindolyl, 3H-indolyl, benzimidazolyl, imidazo[1,2-a]pyridyl, benzothiazolyl, benzoxazolyl, quinoliziny, quinazolinyl, pthalazinyl, quinoxaliny, cinnolinyl, naphthyridinyl, pyrido[3,4-b]pyridyl, pyrido[3,2-b]pyridyl, pyrido[4,3-b]pyridyl, quinolyl, isoquinolyl, 20 tetrazolyl, 5,6,7,8-tetrahydroquinolyl, 5,6,7,8-tetrahydroisoquinolyl, purinyl, pteridinyl, carbazolyl, xanthenyl or benzoquinolyl.

"C₁-C₆-alkyl heteroaryl" refers to C₁-C₆-alkyl groups having a heteroaryl substituent, including 2-furylmethyl, 2-thienylmethyl, 2-(1H-indol-3-yl)ethyl and the like.

"C₂-C₆-alkenyl" refers to alkenyl groups preferably having from 2 to 6 carbon atoms and having at least 1 or 2 sites of alkenyl unsaturation. Preferable alkenyl groups include ethenyl (-CH=CH₂), n-2-propenyl (allyl, -CH₂CH=CH₂) and the like.

5 "C₂-C₆-alkenyl aryl" refers to C₂-C₆-alkenyl groups having an aryl substituent, including 2-phenylvinyl and the like.

"C₂-C₆-alkenyl heteroaryl" refers to C₂-C₆-alkenyl groups having a heteroaryl substituent, including 2-(3-pyridinyl)vinyl and the like.

10 "C₂-C₆-alkynyl" refers to alkynyl groups preferably having from 2 to 6 carbon atoms and having at least 1-2 sites of alkynyl unsaturation, preferred alkynyl groups include ethynyl (-C≡CH), propargyl (-CH₂C≡CH), and the like.

"C₂-C₆-alkynyl aryl" refers to C₂-C₆-alkynyl groups having an aryl substituent, including phenylethynyl and the like.

"C₂-C₆-alkynyl heteroaryl" refers to C₂-C₆-alkynyl groups having a heteroaryl substituent, including 2-thienylethynyl and the like.

15 "C₃-C₈-cycloalkyl" refers to a saturated carbocyclic group of from 3 to 8 carbon atoms having a single ring (e.g., cyclohexyl) or multiple condensed rings (e.g., norbornyl). Preferred cycloalkyl include cyclopentyl, cyclohexyl, norbornyl and the like.

20 "Heterocycloalkyl" refers to a C₃-C₈-cycloalkyl group according to the definition above, in which up to 3 carbon atoms are replaced by heteroatoms chosen from the group consisting of O, S, NR, R being defined as hydrogen or methyl. Preferred heterocycloalkyl include pyrrolidine, piperidine, piperazine, 1-methylpiperazine, morpholine, and the like.

"C₁-C₆-alkyl cycloalkyl" refers to C₁-C₆-alkyl groups having a cycloalkyl substituent, including cyclohexylmethyl, cyclopentylpropyl, and the like.

"C₁-C₆-alkyl heterocycloalkyl" refers to C₁-C₆-alkyl groups having a heterocycloalkyl substituent, including 2-(1-pyrrolidinyl)ethyl, 4-morpholinylmethyl, (1-methyl-4-piperidinyl)methyl and the like.

"Carboxy" refers to the group -C(O)OH.

- 5 "C₁-C₆-alkyl carboxy" refers to C₁-C₆-alkyl groups having a carboxy substituent, including 2-carboxyethyl and the like.

"Acyl" refers to the group -C(O)R where R includes "C₁-C₆-alkyl", "aryl", "heteroaryl", "C₁-C₆-alkyl aryl" or "C₁-C₆-alkyl heteroaryl".

- 10 "C₁-C₆-alkyl acyl" refers to C₁-C₆-alkyl groups having an acyl substituent, including 2-acetyethyl and the like.

"Aryl acyl" refers to aryl groups having an acyl substituent, including 2-acetylphenyl and the like.

"Heteroaryl acyl" refers to heteroaryl groups having an acyl substituent, including 2-acetylpyridyl and the like.

- 15 "C₃-C₈-(hetero)cycloalkyl acyl" refers to 3 to 8 membered cycloalkyl or heterocycloalkyl groups having an acyl substituent.

- 20 "Acyloxy" refers to the group -OC(O)R where R includes H, "C₁-C₆-alkyl", "C₂-C₆-alkenyl", "C₂-C₆-alkynyl", "C₃-C₈-cycloalkyl", heterocycloalkyl "heterocycloalkyl", "aryl", "heteroaryl", "C₁-C₆-alkyl aryl" or "C₁-C₆-alkyl heteroaryl", "C₂-C₆-alkenyl aryl", "C₂-C₆-alkenyl heteroaryl", "C₂-C₆-alkynyl aryl", "C₂-C₆-alkynyl heteroaryl", "C₁-C₆-alkyl cycloalkyl", "C₁-C₆-alkyl heterocycloalkyl".

"C₁-C₆-alkyl acyloxy" refers to C₁-C₆-alkyl groups having an acyloxy substituent, including 2-(acyloxy)ethyl and the like.

"Alkoxy" refers to the group $-O-R$ where R includes "C₁-C₆-alkyl" or "aryl" or "hetero-aryl" or "C₁-C₆-alkyl aryl" or "C₁-C₆-alkyl heteroaryl". Preferred alkoxy groups include by way of example, methoxy, ethoxy, phenoxy and the like.

- 5 "C₁-C₆-alkyl alkoxy" refers to C₁-C₆-alkyl groups having an alkoxy substituent, including 2-ethoxyethyl and the like.

"Alkoxy carbonyl" refers to the group $-C(O)OR$ where R includes H, "C₁-C₆-alkyl" or "aryl" or "heteroaryl" or "C₁-C₆-alkyl aryl" or "C₁-C₆-alkyl heteroaryl".

"C₁-C₆-alkyl alkoxy carbonyl" refers to C₁-C₆-alkyl groups having an alkoxy carbonyl substituent, including 2-(benzyloxy carbonyl)ethyl and the like.

- 10 "Aminocarbonyl" refers to the group $-C(O)NRR'$ where each R, R' includes independently hydrogen or C₁-C₆-alkyl or aryl or heteroaryl or "C₁-C₆-alkyl aryl" or "C₁-C₆-alkyl heteroaryl".

"C₁-C₆-alkyl aminocarbonyl" refers to C₁-C₆-alkyl groups having an aminocarbonyl substituent, including 2-(dimethylaminocarbonyl)ethyl and the like.

- 15 "Acylamino" refers to the group $-NRC(O)R'$ where each R, R' is independently hydrogen, "C₁-C₆-alkyl", "C₂-C₆-alkenyl", "C₂-C₆-alkynyl", "C₃-C₈-cycloalkyl", "heterocycloalkyl", "aryl", "heteroaryl", "C₁-C₆-alkyl aryl" or "C₁-C₆-alkyl heteroaryl", "C₂-C₆-alkenyl aryl", "C₂-C₆-alkenyl heteroaryl", "C₂-C₆-alkynyl aryl", "C₂-C₆-alkynyl heteroaryl", "C₁-C₆-alkyl cycloalkyl", "C₁-C₆-alkyl heterocycloalkyl".

- 20 "C₁-C₆-alkyl acylamino" refers to C₁-C₆-alkyl groups having an acylamino substituent, including 2-(propionylamino)ethyl and the like.

"Ureido" refers to the group $-NRC(O)NR'R''$ where each R, R', R'' is independently hydrogen, "C₁-C₆-alkyl", "C₂-C₆-alkenyl", "C₂-C₆-alkynyl", "C₃-C₈-cycloalkyl", "heterocycloalkyl", "aryl", "heteroaryl", "C₁-C₆-alkyl aryl" or "C₁-C₆-alkyl heteroaryl",

"C₂-C₆-alkenyl aryl", "C₂-C₆-alkenyl heteroaryl", "C₂-C₆-alkynyl aryl", "C₂-C₆-alkynylheteroaryl", "C₁-C₆-alkyl cycloalkyl", "C₁-C₆-alkyl heterocycloalkyl", and where R' and R", together with the nitrogen atom to which they are attached, can optionally form a 3-8-membered heterocycloalkyl ring.

- 5 "C₁-C₆-alkyl ureido" refers to C₁-C₆-alkyl groups having an ureido substituent, including 2-(N'-methylureido)ethyl and the like.

"Carbamate" refers to the group -NRC(O)OR' where each R, R' is independently hydrogen, "C₁-C₆-alkyl", "C₂-C₆-alkenyl", "C₂-C₆-alkynyl", "C₃-C₈-cycloalkyl", "heterocycloalkyl", "aryl", "heteroaryl", "C₁-C₆-alkyl aryl" or "C₁-C₆-alkyl heteroaryl",
 10 "C₂-C₆-alkenyl aryl", "C₂-C₆-alkenyl heteroaryl", "C₂-C₆-alkynyl aryl", "C₂-C₆-alkynylheteroaryl", "C₁-C₆-alkyl cycloalkyl", "C₁-C₆-alkyl heterocycloalkyl".

"Amino" refers to the group -NRR' where each R, R' is independently hydrogen or "C₁-C₆-alkyl" or "aryl" or "heteroaryl" or "C₁-C₆-alkyl aryl" or "C₁-C₆-alkyl heteroaryl", or "cycloalkyl", or "heterocycloalkyl", and where R and R', together with the nitrogen atom to
 15 which they are attached, can optionally form a 3-8-membered heterocycloalkyl ring.

"C₁-C₆-alkyl amino" refers to C₁-C₆-alkyl groups having an amino substituent, including 2-(1-pyrrolidinyl)ethyl and the like.

"Ammonium" refers to a positively charged group -N⁺RR'R'', where each R, R', R'' is independently "C₁-C₆-alkyl" or "C₁-C₆-alkyl aryl" or "C₁-C₆-alkyl heteroaryl", or
 20 "cycloalkyl", or "heterocycloalkyl", and where R and R', together with the nitrogen atom to which they are attached, can optionally form a 3-8-membered heterocycloalkyl ring.

"C₁-C₆-alkyl ammonium" refers to C₁-C₆-alkyl groups having an ammonium substituent, including 2-(1-pyrrolidinyl)ethyl and the like.

"Halogen" refers to fluoro, chloro, bromo and iodo atoms.

- "Sulfonyloxy" refers to a group $-\text{OSO}_2\text{-R}$ wherein R is selected from H, "C₁-C₆-alkyl", "C₁-C₆-alkyl" substituted with halogens, e.g., an $-\text{OSO}_2\text{-CF}_3$ group, "C₂-C₆-alkenyl", "C₂-C₆-alkynyl", "C₃-C₆-cycloalkyl", "heterocycloalkyl", "aryl", "heteroaryl", "C₁-C₆-alkyl aryl" or "C₁-C₆-alkyl heteroaryl", "C₂-C₆-alkenyl aryl", "C₂-C₆-alkenyl heteroaryl", "C₂-C₆-alkynyl aryl", "C₂-C₆-alkynylheteroaryl", "C₁-C₆-alkyl cycloalkyl", "C₁-C₆-alkyl heterocycloalkyl".

"C₁-C₆-alkyl sulfonyloxy" refers to C₁-C₅-alkyl groups having a sulfonyloxy substituent, including 2-(methylsulfonyloxy)ethyl and the like.

- "Sulfonyl" refers to group $-\text{SO}_2\text{-R}$ wherein R is selected from H, "aryl", "heteroaryl", "C₁-C₆-alkyl", "C₁-C₆-alkyl" substituted with halogens, e.g., an $-\text{SO}_2\text{-CF}_3$ group, "C₂-C₆-alkenyl", "C₂-C₆-alkynyl", "C₃-C₆-cycloalkyl", "heterocycloalkyl", "aryl", "heteroaryl", "C₁-C₆-alkyl aryl" or "C₁-C₆-alkyl heteroaryl", "C₂-C₆-alkenyl aryl", "C₂-C₆-alkenyl heteroaryl", "C₂-C₆-alkynyl aryl", "C₂-C₆-alkynylheteroaryl", "C₁-C₆-alkyl cycloalkyl", "C₁-C₆-alkyl heterocycloalkyl".
- "C₁-C₆-alkyl sulfonyl" refers to C₁-C₅-alkyl groups having a sulfonyl substituent, including 2-(methylsulfonyl)ethyl and the like.

- "Sulfinyl" refers to a group $-\text{S(O)-R}$ wherein R is selected from H, "C₁-C₆-alkyl", "C₁-C₆-alkyl" substituted with halogens, e.g., a $-\text{SO-CF}_3$ group, "C₂-C₆-alkenyl", "C₂-C₆-alkynyl", "C₃-C₆-cycloalkyl", "heterocycloalkyl", "aryl", "heteroaryl", "C₁-C₆-alkyl aryl" or "C₁-C₆-alkyl heteroaryl", "C₂-C₆-alkenyl aryl", "C₂-C₆-alkenyl heteroaryl", "C₂-C₆-alkynyl aryl", "C₂-C₆-alkynylheteroaryl", "C₁-C₆-alkyl cycloalkyl", "C₁-C₆-alkyl heterocycloalkyl".

"C₁-C₆-alkyl sulfinyl" refers to C₁-C₅-alkyl groups having a sulfinyl substituent, including 2-(methylsulfinyl)ethyl and the like.

"Sulfanyl" refers to groups $-S-R$ where R includes H , " C_1-C_6 -alkyl", " C_1-C_6 -alkyl" substituted with halogens, e.g., a $-SO-CF_3$ group, " C_2-C_6 -alkenyl", " C_2-C_6 -alkynyl", " C_3-C_6 -cycloalkyl", "heterocycloalkyl", "aryl", "heteroaryl", " C_1-C_6 -alkyl aryl" or " C_1-C_6 -alkyl heteroaryl", " C_2-C_6 -alkenyl aryl", " C_2-C_6 -alkenyl heteroaryl", " C_2-C_6 -alkynyl aryl", " C_2-C_6 -alkynylheteroaryl", " C_1-C_6 -alkyl cycloalkyl", " C_1-C_6 -alkyl heterocycloalkyl". Preferred sulfanyl groups include methylsulfanyl, ethylsulfanyl, and the like.

" C_1-C_6 -alkyl sulfanyl" refers to C_1-C_6 -alkyl groups having a sulfanyl substituent, including 2-(ethylsulfanyl)ethyl and the like.

"Sulfonylamino" refers to a group $-NRSO_2-R'$ where each R, R' includes independently hydrogen, " C_1-C_6 -alkyl", " C_2-C_6 -alkenyl", " C_2-C_6 -alkynyl", " C_3-C_6 -cycloalkyl", "heterocycloalkyl", "aryl", "heteroaryl", " C_1-C_6 -alkyl aryl" or " C_1-C_6 -alkyl heteroaryl", " C_2-C_6 -alkenyl aryl", " C_2-C_6 -alkenyl heteroaryl", " C_2-C_6 -alkynyl aryl", " C_2-C_6 -alkynylheteroaryl", " C_1-C_6 -alkyl cycloalkyl", " C_1-C_6 -alkyl heterocycloalkyl".

" C_1-C_6 -alkyl sulfonylamino" refers to C_1-C_6 -alkyl groups having a sulfonylamino substituent, including 2-(ethylsulfonylamino)ethyl and the like.

"Aminosulfonyl" refers to a group $-SO_2-NRR'$ where each R, R' includes independently hydrogen, " C_1-C_6 -alkyl", " C_2-C_6 -alkenyl", " C_2-C_6 -alkynyl", " C_3-C_6 -cycloalkyl", "heterocycloalkyl", "aryl", "heteroaryl", " C_1-C_6 -alkyl aryl" or " C_1-C_6 -alkyl heteroaryl", " C_2-C_6 -alkenyl aryl", " C_2-C_6 -alkenyl heteroaryl", " C_2-C_6 -alkynyl aryl", " C_2-C_6 -alkynylheteroaryl", " C_1-C_6 -alkyl cycloalkyl", " C_1-C_6 -alkyl heterocycloalkyl".

" C_1-C_6 -alkyl aminosulfonyl" refers to C_1-C_6 -alkyl groups having an aminosulfonyl substituent, including 2-(cyclohexylaminosulfonyl)ethyl and the like.

"Substituted or unsubstituted": Unless otherwise constrained by the definition of the individual substituent, the above set out groups, like "alkyl", "alkenyl", "alkynyl", "aryl" and "heteroaryl" etc. groups can optionally be substituted with from 1 to 5 substituents selected

from the group consisting of "C₁-C₆-alkyl", "C₂-C₆-alkenyl", "C₂-C₆-alkynyl", "cycloalkyl", "heterocycloalkyl", "C₁-C₆-alkyl aryl", "C₁-C₆-alkyl heteroaryl", "C₁-C₆-alkyl cycloalkyl", "C₁-C₆-alkyl heterocycloalkyl", "amino", "ammonium", "acyl", "acyloxy", "acylamino", "aminocarbonyl", "alkoxycarbonyl", "ureido", "aryl",

5 "carbamate", "heteroaryl", "sulfinyl", "sulfonyl", "alkoxy", "sulfanyl", "halogen", "carboxy", trihalomethyl, cyano, hydroxy, mercapto, nitro, and the like. Alternatively said substitution could also comprise situations where neighbouring substituents have undergone ring closure, notably when vicinal functional substituents are involved, thus forming, e.g., lactams, lactones, cyclic anhydrides, but also acetals, thioacetals, amins

10 formed by ring closure for instance in an effort to obtain a protective group.

"Pharmaceutically acceptable cationic salts or complexes" is intended to define such salts as the alkali metal salts, (e.g. sodium and potassium), alkaline earth metal salts (e.g. calcium or magnesium), aluminium salts, ammonium salts and salts with organic amines such as with methylamine, dimethylamine, trimethylamine, ethylamine, triethylamine,

15 morpholine, N-Me-D-glucamine, N,N'-bis(phenylmethyl)-1,2-ethanediamine, ethanolamine, diethanolamine, ethylenediamine, N-methylmorpholine, piperidine, benzathine (N,N'-dibenzylethylenediamine), choline, ethylene-diamine, meglumine (N-methylglucamine), benethamine (N-benzylphenethylamine), diethylamine, piperazine, thromethamine (2-amino-2-hydroxymethyl-1,3-propanediol), procaine as well as amines of

20 formula -NR,R',R'' wherein R, R', R'' is independently hydrogen, alkyl or benzyl. Especially preferred salts are sodium and potassium salts.

"Pharmaceutically acceptable salts or complexes" refers to salts or complexes of the below-identified compounds of formula (I) that retain the desired biological activity. Examples of such salts include, but are not restricted to acid addition salts formed with inorganic acids

25 (e.g., hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid, and the like), and salts formed with organic acids such as acetic acid, oxalic acid, tartaric acid, succinic acid, malic acid, fumaric acid, maleic acid, ascorbic acid, benzoic acid, tannic acid,

pamoic acid, alginic acid, polyglutamic acid, naphthalene sulfonic acid, naphthalene disulfonic acid, and poly-galacturonic acid. Said compounds can also be administered as pharmaceutically acceptable quaternary salts known by a person skilled in the art, which specifically include the quaternary ammonium salt of the formula $-NR_1R'_1R''^+Z^-$, wherein

5 R_1, R'_1, R'' is independently hydrogen, alkyl, or benzyl, C_1-C_6 -alkyl, C_2-C_6 -alkenyl, C_2-C_6 -alkynyl, C_1-C_6 -alkyl aryl, C_1-C_6 -alkyl heteroaryl, cycloalkyl, heterocycloalkyl, and Z is a counterion, including chloride, bromide, iodide, -O-alkyl, toluenesulfonate, methylsulfonate, sulfonate, phosphate, or carboxylate (such as benzoate, succinate, acetate, glycolate, maleate, malate, fumarate, citrate, tartrate, ascorbate, cinnamate, mandelate, and diphenylacetate).

10

"Pharmaceutically active derivative" refers to any compound that upon administration to the recipient, is capable of providing directly or indirectly, the activity disclosed herein.

"Enantiomeric excess" (ee) refers to the products that are obtained by an asymmetric synthesis, i.e. a synthesis involving non-racemic starting materials and/or reagents or a synthesis comprising at least one enantioselective step, whereby a surplus of one enantiomer in

15 the order of at least about 52% ee is yielded.


It has now been found that compounds of the present invention are modulators of the Phosphatoinositides 3-kinases (PI3Ks), particularly of Phosphatoinositides 3-kinase γ (PI3K γ). When the phosphatoinositides 3-kinase (PI3K) enzyme is inhibited by the

20 compounds of the present invention, PI3K is unable to exert its enzymatic, biological and/or pharmacological effects. The compounds of the present invention are therefore useful in the treatment and prevention of autoimmune disorders and/or inflammatory diseases, cardiovascular diseases, neurodegenerative diseases, bacterial or viral infections, kidney diseases, platelet aggregation, cancer, transplantation, graft rejection or lung

25 injuries.

General formula (I) according to the present invention also comprises its tautomers, its geometrical isomers, its optically active forms as enantiomers, diastereomers and its racemate forms, as well as pharmaceutically acceptable salts thereof. Preferred pharmaceutically acceptable salts of the formula (I) are acid addition salts formed with pharmaceutically acceptable acids like hydrochloride, hydrobromide, sulfate or bisulfate, phosphate or hydrogen phosphate, acetate, benzoate, succinate, fumarate, maleate, lactate, citrate, tartrate, gluconate, methanesulfonate, benzenesulfonate, and *para*-toluenesulfonate salts.

A first aspect of the present invention consists in novel compounds of formula (I) :



A is an unsubstituted or substituted 5-8 membered heterocyclic group or an unsubstituted or substituted carbocyclic group. Preferably, A is a heterocyclic moiety.

Said carbocyclic group may be fused with an unsubstituted or substituted aryl, an unsubstituted or substituted heteroaryl, an unsubstituted or substituted cycloalkyl or an unsubstituted or substituted heterocycloalkyl.

Such heterocyclic or carbocyclic groups comprise aryl, heteroaryl, cycloalkyl and heterocycloalkyl, including phenyl, phenantrenyl, cyclopentyl, cyclohexyl, norbornyl, pyrrolidine, piperidine, piperazine, 1-methylpiperazine, morpholine, pyrrolyl, furanyl, thienyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,3,4-triazinyl, 1,2,3-triazinyl, benzofuryl, [2,3-dihydro]benzofuryl,

- isobenzofuryl, benzothienyl, benzotriazolyl, isobenzothienyl, indolyl, isoindolyl, 3H-indolyl, benzimidazolyl, imidazo[1,2-a]pyridyl, benzothiazolyl, benzoxazolyl, quinoliziny, quinazoliny, phthalaziny, quinoxaliny, cinnoliny, naphthyridiny, pyrido[3,4-b]pyridyl, pyrido[3,2-b]pyridyl, pyrido[4,3-b]pyridyl, quinolyl, isoquinolyl, tetrazolyl, 5,6,7,8-tetrahydroquinolyl, 5,6,7,8-tetrahydroisoquinolyl, purinyl, pteridinyl, carbazolyl, xanthenyl or benzoquinolyl

- Further exemplary heterocyclic or carbocyclic groups A include unsubstituted or substituted dioxolenyl, unsubstituted or substituted dioxinyl, unsubstituted or substituted dihydrofuranyl, unsubstituted or substituted (dihydro) furanyl, unsubstituted or substituted (dihydro)oxazinyl, unsubstituted or substituted oxazinoyl, unsubstituted or substituted pyridinyl, unsubstituted or substituted isooxazolyl, unsubstituted or substituted oxazolyl, unsubstituted or substituted (dihydro)naphthalenyl, unsubstituted or substituted pyrimidinyl, unsubstituted or substituted triazolyl, unsubstituted or substituted imidazolyl, unsubstituted or substituted pyrazinyl, unsubstituted or substituted thiazolyl, unsubstituted or substituted thiadiazolyl, unsubstituted or substituted oxadiazolyl.

In one embodiment of the present invention A is a dioxolenyl or a pyridinyl moiety.

X is S, O or $-NR^3$, preferably S. R^3 is selected from the group comprising or consisting of H or C_1-C_6 -alkyl.

Y is S or O, preferably O.

- R^1 is selected from the group comprising or consisting of H, CN, carboxy, acyl, C_1-C_6 -alkoxy, halogen, hydroxy, acyloxy, an unsubstituted or substituted C_1-C_6 -alkyl carboxy, an unsubstituted or substituted C_1-C_6 -alkyl acyloxy, an unsubstituted or substituted C_1-C_6 -alkyl alkoxy, alkoxycarbonyl, an unsubstituted or substituted C_1-C_6 -alkyl alkoxycarbonyl, aminocarbonyl, an unsubstituted or substituted C_1-C_6 -alkyl aminocarbonyl, acylamino, an unsubstituted or substituted C_1-C_6 -alkyl acylamino, ureido, an unsubstituted or substituted C_1-C_6 -alkyl ureido, amino, an unsubstituted or substituted C_1-C_6 -alkyl amino, ammonium,

sulfonyloxy, an unsubstituted or substituted C₁-C₆-alkyl sulfonyloxy, sulfonyl, an unsubstituted or substituted C₁-C₆-alkyl sulfonyl, sulfinyl, an unsubstituted or substituted C₁-C₆-alkyl sulfinyl, sulfanyl, an unsubstituted or substituted C₁-C₆-alkyl sulfanyl, sulfonylamino, an unsubstituted or substituted C₁-C₆-alkyl sulfonylamino or carbamate.

5 Preferably R¹ is H.

R² is selected from the group comprising or consisting of H, halogen, acyl, amino, an unsubstituted or substituted C₁-C₆-alkyl, an unsubstituted or substituted C₂-C₆-alkenyl, an unsubstituted or substituted C₂-C₆-alkynyl, an unsubstituted or substituted C₁-C₆-alkyl carboxy, an unsubstituted or substituted C₁-C₆-alkyl acyl, an unsubstituted or substituted C₁-C₆-alkyl alkoxycarbonyl, an unsubstituted or substituted C₁-C₆-alkyl aminocarbonyl, an unsubstituted or substituted C₁-C₆-alkyl acyloxy, an unsubstituted or substituted C₁-C₆-alkyl acylamino, an unsubstituted or substituted C₁-C₆-alkyl ureido, an unsubstituted or substituted C₁-C₆-alkyl carbamate, an unsubstituted or substituted C₁-C₆-alkyl amino, an unsubstituted or substituted C₁-C₆-alkyl alkoxy, an unsubstituted or substituted C₁-C₆-alkyl sulfanyl, an unsubstituted or substituted C₁-C₆-alkyl sulfinyl, an unsubstituted or substituted C₁-C₆-alkyl sulfonyl, an unsubstituted or substituted C₁-C₆-alkyl sulfonylaminoaryl, aryl, heteroaryl, an unsubstituted or substituted C₃-C₆-cycloalkyl or heterocycloalkyl, an unsubstituted or substituted C₁-C₆-alkyl aryl, an unsubstituted or substituted C₁-C₆-alkyl heteroaryl, an unsubstituted or substituted C₂-C₆-alkenyl-aryl or -heteroaryl, an unsubstituted or substituted C₂-C₆-alkynyl aryl or -heteroaryl, carboxy, cyano, hydroxy, C₁-C₆-alkoxy, nitro, acylamino, ureido, sulfonylamino, sulfanyl, or sulfonyl. Preferably R² is H.

10
15
20

In a specific embodiment, R¹ and R² are both H.


G is a substituted or unsubstituted C₁-C₆-alkyl, substituted or unsubstituted C₂-C₆-alkenyl, substituted or unsubstituted C₂-C₆-alkynyl, substituted or unsubstituted heteroaryl, an unsubstituted or substituted C₁-C₆-alkyl aryl, an unsubstituted or substituted C₁-C₆-alkyl heteroaryl, an unsubstituted or substituted C₂-C₆-alkenyl-aryl or -heteroaryl, an

25

unsubstituted or substituted C_2-C_6 -alkynyl aryl or -heteroaryl, substituted or unsubstituted C_1-C_6 -alkoxy, cyano, substituted or unsubstituted C_1-C_6 -acyl or G is a sulfonyl moiety.

In particular, G is selected from the group comprising or consisting of a sulfonyl moiety, a cyano or an substituted or unsubstituted C_1-C_6 -alkoxy.

- 5 In a chemical library 4 compounds of formula (I) are disclosed :



The compounds are tautomers of formula (I). No biological activity is disclosed for said 4 compounds.

- In one embodiment of the present invention G is a sulfonyl moiety of the formula $-SO_2-R^4$,
 10 whereby R^4 is selected from the group comprising or consisting of H, unsubstituted or substituted C_1-C_6 -alkyl, unsubstituted or substituted C_2-C_6 -alkenyl, unsubstituted or substituted C_2-C_6 -alkynyl, unsubstituted or substituted C_1-C_6 -alkyl carboxy, an unsubstituted or substituted C_1-C_6 -alkyl acyl, an unsubstituted or substituted C_1-C_6 -alkyl alkoxycarbonyl, an unsubstituted or substituted C_1-C_6 -alkyl aminocarbonyl, an
 15 unsubstituted or substituted C_1-C_6 -alkyl acyloxy, an unsubstituted or substituted C_1-C_6 -alkyl acylamino, an unsubstituted or substituted C_1-C_6 -alkyl ureido, an unsubstituted or substituted C_1-C_6 -alkyl carbamate, an unsubstituted or substituted C_1-C_6 -alkyl amino, an unsubstituted or substituted C_1-C_6 -alkoxy, an unsubstituted or substituted C_1-C_6 -alkyl

sulfanyl, an unsubstituted or substituted C₁-C₆-alkyl sulfinyl, an unsubstituted or substituted C₁-C₆-alkyl sulfonyl, an unsubstituted or substituted C₁-C₆-alkyl sulfonylaminoaryl, aryl, heteroaryl, an unsubstituted or substituted C₃-C₈-cycloalkyl or heterocycloalkyl, an unsubstituted or substituted C₁-C₆-alkyl aryl, an unsubstituted or substituted C₁-C₆-alkyl heteroaryl, an unsubstituted or substituted C₂-C₆-alkenyl-aryl or -heteroaryl, an unsubstituted or substituted C₂-C₆-alkynyl aryl or -heteroaryl, carboxy, hydroxy, C₁-C₆-alkoxy, acylamino, sulfonylamino.

In one embodiment of the present invention R⁴ is an unsubstituted or substituted aryl, an unsubstituted or substituted heteroaryl, an unsubstituted or substituted C₁-C₃ alkyl.

In a specific embodiment, X is S, Y is O, R¹ and R² are H, A is a dioxolenyl or a pyridinyl moiety.

The compounds according to formula (I) are suitable for the modulation, notably the inhibition of the activity of phosphatoinositides 3-kinases (PI3K), particularly phosphatoinositides 3-kinase (PI3Kγ). It is therefore believed that the compounds of the present invention are also particularly useful for the treatment and/or prevention of disorders which are mediated by PI3Ks, particularly PI3Kγ. Said treatment involves the modulation – notably the inhibition or the down regulation – of the phosphatoinositides 3-kinases.

The compounds of the present invention may be obtained as E/Z isomer mixture or as essentially pure E-isomers or Z isomers. The E/Z isomerism preferably refers to the vinyl moiety linking the phenyl with the azolidinone moiety. In a specific embodiment, the compounds of formula (I) are Z-isomers.

Compounds of the present invention include in particular those of the group consisting of:

Example	Name
1	N-(5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-2-chloro-benzenesulfonamide
2	Ethanesulfonic acid (5-benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-amide
3	N-(5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-3-chloro-benzenesulfonamide
4	5-Chloro-1,3-dimethyl-1H-pyrazole-4-sulfonic acid (5-benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-amide
5	3-(5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidenesulfamoyl)-thiophene-2-carboxylic acid methyl ester
6	Preparation of 6-Chloro-pyridine-3-sulfonic acid (5-benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-amide
7	Quinoline-8-sulfonic acid (5-benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-amide
8	N-(5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-benzenesulfonamide
9	N-(5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-4-methyl-benzenesulfonamide
10	N-(5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-methanesulfonamide
11	N-[5-(2,2-Difluoro-benzo[1,3]dioxol-5-ylmethylene)-4-oxo-thiazolidin-2-ylidene]-benzenesulfonamide
12	N-[5-(2,2-Difluoro-benzo[1,3]dioxol-5-ylmethylene)-4-oxo-thiazolidin-2-ylidene]-4-methyl-benzenesulfonamide
13	N-[5-(2,2-Difluoro-benzo[1,3]dioxol-5-ylmethylene)-4-oxo-thiazolidin-2-ylidene]-methanesulfonamide

- 14 Biphenyl-2-sulfonic acid (5-benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-amide
- 15 Pyridine-3-sulfonic acid (5-benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-amide
- 16 3-(4-Oxo-5-quinolin-6-ylmethylene-thiazolidin-2-ylidenesulfamoyl)-thiophene-2-carboxylic acid methyl ester
- 17 2-Chloro-N-(4-oxo-5-quinolin-6-ylmethylene-thiazolidin-2-ylidene)-benzenesulfonamide
- 18 3-(5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidenesulfamoyl)-thiophene-2-carboxylic acid
- 19 5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene-cyanamide
- 20 5-Benzo[1,3]dioxol-5-ylmethylene-thiazolidine-2,4-dione 2-(O-methyl-oxime)
- 21 Preparation of 4-Oxo-5-quinoxalin-6-ylmethylene-thiazolidin-2-ylidene-cyanamide
- 22 5-Benzo[1,3]dioxol-5-ylmethylene-2-benzylimino-thiazolidin-4-one
- 23 2-Benzylimino-5-quinolin-6-ylmethylene-thiazolidin-4-one
- 24 2-Propylimino-5-quinolin-6-ylmethylene-thiazolidin-4-one
- 25 5-Benzo[1,3]dioxol-5-ylmethylene-2-propylimino-thiazolidin-4-one
- 26 5-(4-Dimethylamino-quinazolin-6-ylmethylene)-2-methylamino-thiazol-4-one

The compounds of the present invention are useful as medicaments. They may be used for the preparation of a medicament for the prophylaxis and/or treatment of autoimmune disorders and/or inflammatory diseases, cardiovascular diseases, neurodegenerative

diseases, bacterial or viral infections, kidney diseases, platelet aggregation, cancer, transplantation, graft rejection or lung injuries.

5 In one embodiment, the compounds of formula (I) are useful for the treatment and/or prophylaxis of autoimmune diseases or inflammatory diseases such as multiple sclerosis, psoriasis, rheumatoid arthritis, systemic lupus erythematosus, inflammatory bowel disease, lung inflammation, thrombosis or brain infection/inflammation such as meningitis or encephalitis.

10 In another embodiment, the compounds of formula (I) are useful for the treatment and/or prophylaxis of neurodegenerative diseases including multiple sclerosis, Alzheimer's disease, Huntington's disease, CNS trauma, stroke or ischemic conditions.

In still a further embodiment according to the invention, the compounds of formula (I) are useful for the treatment and/or prophylaxis of cardiovascular diseases such as atherosclerosis, heart hypertrophy, cardiac myocyte dysfunction, elevated blood pressure or vasoconstriction.

15 In still another embodiment according to the invention, the compounds of formula (I) are useful for the treatment and/or prophylaxis of chronic obstructive pulmonary disease, anaphylactic shock fibrosis, psoriasis, allergic diseases, asthma, stroke or ischemic conditions, ischemia-reperfusion, platelets aggregation/activation, skeletal muscle atrophy/hypertrophy, leukocyte recruitment in cancer tissue, angiogenesis, invasion
20 metastasis, in particular melanoma, Kaposi's sarcoma, acute and chronic bacterial and viral infections, sepsis, transplantation, graft rejection, glomerulo sclerosis, glomerulo nephritis, progressive renal fibrosis, endothelial and epithelial injuries in the lung or in general lung airways inflammation.


25 Still a further object of the present invention is a process for preparing 2-imino-azolinone-vinyl fused-benzene derivatives according to formula (I).

The 2-imino-azolinone-vinyl fused-benzene derivatives exemplified in this invention may be prepared from readily available starting materials using the following general methods and procedures. It will be appreciated that where typical or preferred experimental conditions (i.e. reaction temperatures, time, moles of reagents, solvents etc.) are given, other experimental conditions can also be used unless otherwise stated. Optimum reaction conditions may vary with the particular reactants or solvents used, but such conditions can be determined by the person skilled in the art, using routine optimisation procedures.


Methods of preparing the compounds within formula (I).

Generally, the 2-imino-azolinone-vinyl fused-benzene derivatives according to the general formula (I) could be obtained by several synthetic approaches, using both solution-phase and solid-phase chemistry protocols (Brummond et al., *J.O.C.*, 64, 1723-1726 (1999)), either by conventional methods or by microwave-assisted techniques (see schemes 1, 2 and 3). In a first step, approximately equimolar amounts of the reactant P1 and reagent P2 (2-amino-4,5-dihydro-1,3-thiazol-4-one, 2-Imino-thiazolidine-4-thione, 2-Imino-oxazolidin-4-one, 2-Imino-oxazolidine-4-thione, 2-Imino-1-alkyl-imidazolidin-4-one or 2-Imino-1-alkyl-imidazolidine-4-thione) or reagent P3 (Oxazolidine-2,4-dithione, 2-Thioxo-oxazolidin-4-one, 1-Alkyl-2-thioxo-imidazolidin-4-one, 1-Alkyl-imidazolidine-2,4-dithione, Thiazolidine-2,4-dithione or rhodanin) are heated in the presence of a mild base to provide the corresponding olefin of formula (Ia) or (Ib) respectively.

2-imino-azolinone-vinyl fused-benzene derivatives can be obtained by reacting intermediate (Ia) with sulfonylhalides or acylhalides (L-G, L=leaving group) in the presence of a scavenger base affording compounds of formula (I)



In case G is alkyl or aryl, the 2-imino-azolinone-vinyl fused-benzene derivatives of formula (I) can be accessed through the reaction of intermediate (Ib) with the corresponding amines, as set out in Scheme 2.



- While the first step leading to intermediates (Ia) and (Ib) may be carried out in the absence of a solvent at a temperature, which is sufficiently high to cause at least partial melting of the reaction mixture, it is preferably carried out in the presence of a reaction inert solvent. A preferred such temperature is in the range of from 100°C to 250°C, and especially
- 5 preferred is a temperature of from 120°C to 200°C. Examples of such solvents for the above reaction include solvents like dimethoxymethane, xylene, toluene, o-dichlorobenzene etc. Examples of suitable mild bases for the above reaction are alkali metal and alkaline earth salts of weak acids such as the (C₁-C₁₂)-alkyl carboxylic acids and benzoic acid, alkali metal and alkaline earth carbonates and bicarbonates such as calcium carbonate,
- 10 magnesium carbonate, potassium bicarbonate and secondary amines such as piperidine, morpholine as well as tertiary amines such as pyridine, triethylamine, diisopropylethylamine, N-methylmorpholine, N-Ethylpiperidine, N-Methylpiperidine and the like. Especially preferred mild bases are sodium acetate or piperidine for reasons of economy and efficiency.
- 15 In a typical such reaction (Tietze et.al., in "The Knoevenagel reaction", p.341 ff., Pergamon Press, Oxford 1991, Eds.: Trost B.M., Fleming I.) the aldehyde starting material P1 and the corresponding heterocycle P2 (2-amino-4,5-dihydro-1,3-thiazol-4-one, 2-Imino-thiazolidine-4-thione, 2-Imino-oxazolidin-4-one, 2-Imino-oxazolidine-4-thione, 2-Imino-1-alkyl-imidazolidin-4-one or 2-Imino-1-alkyl-imidazolidine-4-thione) or heterocycle P3
- 20 (Oxazolidine-2,4-dithione, 2-Thioxo-oxazolidin-4-one, 1-Alkyl-2-thioxo-imidazolidin-4-one, 1-Alkyl-imidazolidine-2,4-dithione, Thiazolidine-2,4-dithione or rhodanin) are combined in approximately equimolar amounts with 0.5 to one equivalent of piperidine in dimethoxymethane or similar solvent and heated between 120 and 200°C at which the reaction is substantially complete in from 15 minutes to 3 hours. The desired olefins of
- 25 formula (Ia) or (Ib) respectively are then isolated by filtration, in case they precipitated out of the reaction mixture upon cooling, or for example, by mixing with water and subsequent filtration, to obtain the crude products, which are purified, if desired, e.g. by crystallization or by standard chromatographic methods.

Alternatively olefins of formula (Ia) or (Ib) respectively may be obtained typically by mixing equimolar amounts of P2 (2-amino-4,5-dihydro-1,3-thiazol-4-one, 2-Imino-thiazolidine-4-thione, 2-Imino-oxazolidin-4-one, 2-Imino-oxazolidine-4-thione, 2-Imino-1-alkyl-imidazolidin-4-one or 2-Imino-1-alkyl-imidazolidine-4-thione) or heterocycle P3 respectively (Oxazolidine-2,4-dithione, 2-Thioxo-oxazolidin-4-one, 1-Alkyl-2-thioxo-imidazolidin-4-one, 1-Alkyl-imidazolidine-2,4-dithione, Thiazolidine-2,4-dithione or rhodanin) with aldehyde P1 and molar excess, preferably a 2 to 4 fold excess, of anhydrous sodium acetate and the mixture is heated at a temperature high enough to effect melting, at which temperature the reaction is mainly complete in from 5 to 60 minutes.


More preferred reactions conditions are where the above reactions are carried out in acidic media such as acetic acid in the presence of sodium acetate, α -amino-acids or β -alanine. 2-amino-4,5-dihydro-1,3-thiazol-4-one (P2) or rhodanin (P3) are mixed with equimolar amounts of aldehyde P1 in the presence of β -alanine in the range of 0.1 to 1 equivalent in acetic acid. The reaction mixture is heated between 80° to 130°C for 5 minutes to 5 hours, affording the intermediates (Ia) and (Ib) as precipitates. Filtration and washing with water afford compounds in high purity.

Above described reactions can be carried out alternatively under microwave conditions as heating source between 140°C and 240°C at which the reaction is substantially complete from 3 to 10 minutes.

In case G is substituted or un-substituted alkyl- or aryl-sulfonyl group or substituted or un-substituted alkyl- or aryl-carbonyl group conditions, as shown on scheme 1 are applied. Typically intermediate (Ia) is dissolved in an aprotic solvent such as NMP or DMA. This solution is treated with at least one equivalent, preferably two to three equivalents of tertiary amine such as pyridine, triethylamine, diisopropylethylamine, N-methylmorpholine, N-Ethylpiperidine, N-Methylpiperidine and the like. Especially preferred bases are triethylamine or diisopropylethylamine, followed by the addition of the corresponding sulfonyl- or acylchloride at reaction temperatures between 0° to 50°C.

Typically the reaction mixtures are stirred between 0.5 to 15 hours, upon which the solvent is evaporated and the final 2-imino-azolinone-vinyl fused-benzene derivatives are precipitated using water and ethylacetate. Standard chromatography techniques may be applied to reach required purities.

- 5 In case G is a substituted or un-substituted alkyl or aryl-group, the 2-imino-azolinone-vinyl fused-benzene derivatives of formula (I) can be accessed through the reaction of intermediate (Ib) with the corresponding substituted or un-substituted alkyl- or arylamines. Intermediates (Ib) are reacted with 1 to 10 equivalents of the corresponding amines in the presence of an inorganic base in reaction solvents like MeOH, EtOH, Acetonitrile, DME
- 10 and the like. Preferably the solvent is mixed with up to 50% of water. Preferred inorganic bases are K_2CO_3 , $CaCO_3$, Na_2CO_3 , $BaCO_3$ and the like. Typical reaction times are 3 to 15 hours under solvent reflux. Typically 2-imino-azolinone-vinyl fused-benzene derivatives of this type precipitate out of the reaction mixture. In some cases additional water maybe needed for precipitation in order to afford cpds of formula (I) in high purity and quantity.
- 15 In case G is a cyano-group, substituted or un-substituted oxime-ether reaction scheme 3 is applied to afford 2-imino-azolinone-vinyl fused-benzene derivatives. Intermediate (Ib) is methylated affording intermediate (Ic), which is ultimately transformed into compounds of formula (I) using the corresponding carbodiimide.



Scheme 3

Typically methylation of intermediate (Ib) takes place in the presence of base such as pyridine, triethylamine, diisopropylethylamine, N-methylmorpholine, N-ethylpiperidine, N-methylpiperidine and the like and an alkylating agent such methyl iodide, dimethylsulfate or the like in an inert solvent, which remains unaffected by the presence of alkylating agents. The reaction mixture may be stirred at 25°C to 60°C, preferably at room temperature between 0.5 to 15 hours. Most preferred conditions are the use of methyl iodide in the presence of Hünig's base in tetrahydrofuran or dioxane. Excess of reagents may be easily removed after completion by applying a vacuum to the reaction. Compounds of formula (Ic) easily precipitate upon adding of water.

In a typical reaction where the S-alkyl group of compounds of formula (Ic) is replaced by a NH₂-G moiety leading to compounds of formula (I) the intermediates (Ic) are treated with a strong base such potassium-tert.butoxide, potassiumhydride, sodiumhydride, preferably with potassium tert.butoxide in an inert solvent, which remains unaffected by the presence of a strong base. The mixture is subsequently treated with the corresponding nucleophile, such as cyanamide, substituted or unsubstituted oxime-ether, hydroxylamine and heated

between 50 and 150°C, preferably at 80°C for 1 to 15 hours. Excess of reagents may be removed by standard washing procedures, where upon the compounds of formula (I) precipitate.

If the above set of general synthetic methods are not applicable to obtain compounds according to formula (I) and/or to necessary intermediates for the synthesis of compounds of formula (I), suitable methods of preparation known by a person skilled in the art should be used. In general, the synthesis pathways for any individual compound of formula (I) will depend on the specific substituents of each molecule and upon the ready availability of intermediates necessary; again such factors being appreciated by those of ordinary skill in the art. For all the protection and deprotection methods, see Philip J. Kocienski, in "Protecting Groups", Georg Thieme Verlag Stuttgart, New York, 1994 and, Theodora W. Greene and Peter G. M. Wuts in "Protective Groups in Organic Synthesis", Wiley Interscience, 3rd Edition 1999.

Compounds of this invention can be isolated in association with solvent molecules by crystallization from evaporation of an appropriate solvent. The pharmaceutically acceptable acid addition salts of the compounds of formula (I), which contain a basic center, may be prepared in a conventional manner. For example, a solution of the free base may be treated with a suitable acid, either neat or in a suitable solution, and the resulting salt isolated either by filtration or by evaporation under vacuum of the reaction solvent. Pharmaceutically acceptable base addition salts may be obtained in an analogous manner by treating a solution of compound of formula (I) with a suitable base. Both types of salts may be formed or interconverted using ion-exchange resin techniques.

When employed as pharmaceuticals, the compounds of the present invention are typically administered in the form of a pharmaceutical composition. Hence, pharmaceutical compositions comprising a compound of formula (I) and a pharmaceutically acceptable carrier, diluent or excipient therefore are also within the scope of the present invention. A

person skilled in the art is aware of a whole variety of such carrier, diluent or excipient compounds suitable to formulate a pharmaceutical composition.

The compounds of the invention, together with a conventionally employed adjuvant, carrier, diluent or excipient may be placed into the form of pharmaceutical compositions and unit dosages thereof, and in such form may be employed as solids, such as tablets or filled capsules, or liquids such as solutions, suspensions, emulsions, elixirs, or capsules filled with the same, all for oral use, or in the form of sterile injectable solutions for parenteral (including subcutaneous use). Such pharmaceutical compositions and unit dosage forms thereof may comprise ingredients in conventional proportions, with or without additional active compounds or principles, and such unit dosage forms may contain any suitable effective amount of the active ingredient commensurate with the intended daily dosage range to be employed.

Pharmaceutical compositions containing 2-iminoazolinone-vinyl fused-benzenederivatives of this invention can be prepared in a manner well known in the pharmaceutical art and comprise at least one active compound. Generally, the compounds of this invention are administered in a pharmaceutically effective amount. The amount of the compound actually administered will typically be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the like.

The pharmaceutical compositions of the present invention can be administered by a variety of routes including oral, rectal, transdermal, subcutaneous, intravenous, intramuscular and intranasal. The compositions for oral administration can take the form of bulk liquid solutions or suspensions, or bulk powders. More commonly, however, the compositions are presented in unit dosage forms to facilitate accurate dosing. The term "unit dosage forms" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to

produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient. Typical unit dosage forms include prefilled, premeasured ampoules or syringes of the liquid compositions or pills, tablets, capsules or the like in the case of solid compositions. In such compositions, the 2-iminoazolinone-vinyl fused-benzene derivative is usually a minor component (from about 0.1 to about 50% by weight or preferably from about 1 to about 40% by weight) with the remainder being various vehicles or carriers and processing aids helpful for forming the desired dosing form.

Liquid forms suitable for oral administration may include a suitable aqueous or nonaqueous vehicle with buffers, suspending and dispensing agents, colorants, flavors and the like.

Solid forms may include, for example, any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatine; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

Injectable compositions are typically based upon injectable sterile saline or phosphate-buffered saline or other injectable carriers known in the art. As above mentioned, the 2-iminoazolinone-vinyl fused-benzenederivatives derivatives of formula (I) in such compositions is typically a minor component, frequently ranging between 0.05 to 10% by weight with the remainder being the injectable carrier and the like.

The above described components for orally administered or injectable compositions are merely representative. Further materials as well as processing techniques and the like are set out in Part 5 of *Remington's Pharmaceutical Sciences*, 20th Edition, 2000, Marck Publishing Company, Easton, Pennsylvania, which is incorporated herein by reference.

The compounds of this invention can also be administered in sustained release forms or from sustained release drug delivery systems. A description of representative sustained

release materials can also be found in the incorporated materials in *Remington's Pharmaceutical Sciences*.

In the following the present invention shall be illustrated by means of some examples, which are not construed to be viewed as limiting the scope of the invention. The following abbreviations are hereinafter used in the accompanying examples: min (minute), hr (hour), g (gram), mmol (millimole), m.p. (melting point), eq (equivalents), ml (milliliter), μ l (microliters), ACN (acetonitrile), Boc (butoxycarbonyl), Cbz (carboxybenzyl), CDCl_3 (deuterated chloroform), cHex (cyclohexane), dba (dibenzylideneacetone), DCM (dichloromethane), DEAD (diethylazodicarboxylate), DIC (diisopropylcarbodiimide), DIEA (diisopropylethylamine), DMAP (4-dimethylaminopyridine), DMB (dimethoxyethane), DMF (dimethylformamide), DMSO (dimethylsulfoxide), $\text{DMSO-}d_6$ (deuterated dimethylsulfoxide), EDC (1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride), EtOAc (ethylacetate), Et_2O (diethylether), Fmoc (9-fluorenylmethoxy-carbonyl), HOBt (1-hydroxybenzotriazole), K_2CO_3 (potassium carbonate), MgSO_4 (magnesium sulfate), MsCl (methylsulfonylchloride), MTBE (*tert*-butylmethylether), NaH (sodium hydride), NaHCO_3 (sodium bicarbonate), nBuLi (n-butyllithium), PCC (pyridinium chlorochromate), PE (petroleum ether), QCl (tetrabutylammonium chloride), rt (room temperature), TBTU (O-benzotriazolyl- N,N,N',N' -tetramethyluronium-tetrafluoroborate), TEA (triethylamine), TFA (trifluoroacetic acid), THF (tetrahydrofuran), TMOF (trimethylorthoformate), TMAD (N,N,N',N' -tetramethylazodicarboxamide), TosCl (toluenesulfonylchloride).

Examples:

The following intermediate commercially available aldehydes were used:

Piperonal, 6-Quinolincarboxaldehyde, 6-Quinoxalinecarboxaldehyde, 2,2-Difluoro-1,3-benzodioxole-5-carboxaldehyde.

25 The following intermediates were prepared:

Intermediate 1: Preparation of 4-N-dimethylaminoquinazoline-6-carboxaldehyde



Step I: 4-Nitro isophthalic acid

A mixture of 3-methyl-4-nitrobenzoic acid (150g, 0.825mol), pyridine (1.5L) and water (1.5L) was heated to reflux. To the hot reaction mixture was added KMnO_4 (10mol) portion wise and reflux for 72h. The hot reaction mixture was filtered through celite and washed with hot water. The filtrate was concentrated under vacuum, residue diluted with water (750mL) and acidified with con. HCl at 0°C. The solid obtained was filtered, washed with water and dried under vacuum to give 4-nitro isophthalic acid (98g, 56%).
TLC, Chloroform/Methanol, 7:3, R_f =0.2

Step II: 4-Amino isophthalic acid

To a solution of 4-nitro isophthalic acid (98g, 0.457mol) in methanol (5L) was added Pd/C (20%) and hydrogenated at RT for 4h. The reaction mixture was filtered through celite and filtrate concentrated under vacuum to give 4-amino isophthalic acid (72g, 87%) as a solid.
TLC, Chloroform/Methanol, 7:3, R_f =0.4

Step III: 4-Oxo-3,4-dihydroquinazolin-6-carboxylic acid

A mixture of 4-amino isophthalic acid (17g, 0.093mol) and formamide (85mL) was heated at 180°C for 5h. The reaction mixture was cooled to RT and added acetone. The solid precipitate thus obtained was stirred for 2h, filtered and dried to give 4-oxo-3,4-dihydroquinazoline-6-carboxylic acid (11g, 61%).
TLC, Chloroform/Methanol, 8:2, R_f =0.25

Step IV: 4-Oxo-3,4-dihydroquinazoline-6-methyl carboxylate

To a solution of 4-oxo-3,4-dihydroquinazoline-6-carboxylic acid (24g, 0.126mol) in dry methanol (800mL) was added thionylchloride (37g) at 5°C and then refluxed at 80°C for

5h. The reaction mixture was concentrated under vacuum and crude taken in ethylacetate (250mL). The organic layer was washed with 10% aqueous NaHCO_3 , water, brine and dried. The solvent was removed under vacuum to give 4-oxo-3,4-dihydroquinazoline-6-methyl carboxylate (24g, 92%) as a solid.

5 TLC, Chloroform/Methanol, 8:2, $R_f=0.6$

Step V: Methyl-4-chloroquinazoline-6-carboxylate

A mixture of 4-oxo-3,4-dihydroquinazolin-6-methyl carboxylate (12g, 0.058mol) and phosphorylchloride (180mL) was heated to reflux for 7h. Excess phosphorylchloride was distilled off and crude taken in ethyl acetate (250mL). The organic layer was washed with 10% aqueous NaHCO_3 solution, water, brine and dried. The solvent was removed under vacuum and crude purified by column chromatography over silica gel (30% ethylacetate in pet. ether) to give methyl-4-chloroquinazoline-6-carboxylate (4.5g, 34%) as a solid.

TLC, pet. ether/EtOAc, 1:1, $R_f=0.65$

Step VI: 4-Chloroquinazoline-6-yl methanol

To a solution of methyl-4-chloroquinazoline-6-carboxylate (3.5g, 0.015mol) in dry THF (35mL) at -25°C was added DIBAL-H (4.4g, 0.031mol) and stirred at -25°C to RT for 2h. The reaction mixture was cooled to -10°C and quenched with 10% aqueous NaHCO_3 (9mL). The reaction mixture was extracted with ethylacetate (100mL), washed with water, brine and dried. The solvent was removed under vacuum to give 4-chloroquinazoline-6-yl methanol (2g, 66%).

TLC, Chloroform/Methanol, 8:2, $R_f=0.35$

Step VII: 4-Chloroquinazoline-6-carboxaldehyde

To a solution of 4-chloroquinazoline-6-yl methanol (3.5g, 0.018mol) in dry CH_2Cl_2 (100mL) was added Dess-Martin periodinane (8.4g, 0.019mol) and stirred at RT for 30min. The reaction mixture was washed with 10% aqueous NaHCO_3 (75mL), water, brine and

dried. The solvent was removed under vacuum to give 4-chloroquinazoline-6-carboxaldehyde (3g, 88%) as pale yellow solid.

TLC, Chloroform/Methanol, 9:1, R_f =0.6


5 Step VIII: 4-N-dimethylaminoquinazoline-6-carboxaldehyde

In a flask of 100 ml 4-chloroquinazoline-6-carboxaldehyde (200mg, 1mmol) was dissolved in dioxane (15ml). To this solution was added an aqueous solution of dimethylamine (585mg, 5mmol) in 12 ml water, and the yellow mixture was stirred for two hours at room temperature. After evaporating the solvents in vacuo a yellow solid was obtained (190mg, yield:91%) which was used without further purification.

HPLC: 0.82 min. LC-MS: M/Z ESI: 1.02 min, 202.12 (M+1). NMR: ^1H NMR (DMSO- d_6) δ 10.08 (s, 1H), 8.75 (s, 1H), 8.53 (s, 1H), 8.10 (d, J =9Hz, 1H), 7.78 (d, J =9Hz, 1H), 3.41 (s, 6H).

15 Intermediate 2: Preparation of 5-Benzo[1,3]dioxol-5-ylmethylene-2-imino-thiazolidin-4-

one




In a 100ml round bottom flask were placed 3.87g of pseudohydantoin, 5g of piperonal and 1.92 g of beta-alanine in 30ml of acetic acid. The reaction was stirred for 3h at 100°C and then slowly cooled to room temperature, while the desired condensation product crystallized. The crystals were filtered and washed with acetic acid (rt) affording 8.0g of pure 5-Benzo[1,3]dioxol-5-ylmethylene-2-imino-thiazolidin-4-one.

HPLC: 2.29 min. LC-MS: M/Z ESI: 1.24 min, 249.12 (M+1). NMR: ^1H NMR (DMSO- d_6) δ 9.35 (br. s, 1H), 9.09 (br s, 1H), 7.52 (s, 1H), 7.04-7.13 (m, 3H), 6.13 (s, 2H).


The following intermediates were synthesized according to the synthesis of intermediate 2 using suitable starting materials.

Intermediate 3: Preparation of 2-Amino-5-(2,2-difluoro-benzo[1,3]dioxol-5-ylmethylene)-thiazol-4-one




HPLC: 3.02 min. LC-MS: M/Z ESI: 1.53 min, 285.12 (M+1). NMR: ¹H NMR (DMSO-d₆) δ 9.48 (br. s, 1H), 9.22 (br s, 1H), 7.44-7.61 (m, 3H), 7.41 (d, J=3Hz, 1H).

Intermediate 4: Preparation of 2-Imino-5-quinolin-6-ylmethylene-thiazolidin-4-one




HPLC: 1.19 min. LC-MS: M/Z ESI: 1.16 min, 256.14 (M+1). NMR: ¹H NMR (DMSO-d₆) δ 9.50 (br. s, 1H), 9.24 (br s, 1H), 8.94 (dd, J=6.1;1.7Hz, 1H), 8.42 (d, J=7.5Hz, 1H), 8.18 (d, J=1.86Hz, 1H), 8.11 (d, J=8.6Hz, 1H), 7.92 (dd, J=6.1, 1.7Hz, 1H), 7.76 (s, 1H), 7.59 (dd, J=4.1, 8.3Hz, 1H).

Intermediate 5: Preparation of 5-(4-Dimethylamino-quinazolin-6-ylmethylene)-2-thioxo-thiazolidin-4-one.




HPLC: 1.97 min. LC-MS: M/Z ESI: 1.23 min, 317.10 (M+1). NMR: ¹H NMR (DMSO-d₆) δ 14.25 (br. s, 1H), 8.80 (s, 1H), 8.55 (s, 1H), 8.12 (d, J=8.7Hz, 1H), 7.92 (m, 2H).

Intermediate 6: Preparation of 5-Quinolin-6-ylmethylene-2-thioxo-thiazolidin-4-one




5 HPLC: 2.11 min. LC-MS: M/Z ESI: 1.23 min, 273.10 (M+1). NMR: ¹H NMR (DMSO-d₆)
 δ 13.9 (s, b 1H), 8.97 (dd, J=1.9Hz, 4.1Hz, 1H), 8.52 (d, J=7.9Hz, 1H), 8.23 (s, 1H), 8.11
 (d, J=9.0Hz, 1H), 7.96 (dd, J=1.9Hz, 4.1Hz, 1H), 7.79 (s, 1H), 7.61 (dd, J=4.1Hz, 8.3Hz,
 1H)

10 Intermediate 7: Preparation of 5-Benzo[1,3]dioxol-5-ylmethylene-2-thioxo-thiazolidin-4-one




HPLC: 3.55 min. LC-MS: M/Z ESI: 1.33 min, 266.12 (M-1). NMR: ¹H NMR (DMSO-d₆)
 δ 12.5 (br. s, 1H), 7.73 (s, 1H), 7.06-7.18 (m, 3H), 6.05 (s, 2H).

Intermediate 8: Preparation of 5-Quinoxaline-6-ylmethylene-2-thioxo-thiazolidin-4-one



15

HPLC: 3.01 min. LC-MS: M/Z ESI: 1.17 min, 272.10 (M-1). NMR: ¹H NMR (DMSO-d₆)
 δ 14.0 (br. s, 1H), 9.00 (s, 2H), 8.31 (s, 1H), 8.21 (d, J=8.7Hz, 1H), 8.05 (d, J=8.7Hz, 1H),
 7.90 (s, 1H).

Intermediate 9: 5-Benzo[1,3]dioxol-5-ylmethylene-2-methylsulfanyl-thiazol-4-one


2g (7.54mmol) of 5-Benzo[1,3]dioxol-5-ylmethylene-2-thioxo-thiazolidin-4-one and 1.5 ml (1.15 eq.) of DIEA were dissolved in 80ml NMP. To this solution was added dropwise a freshly prepared solution of 2.43ml (5 eq.) of Methyl iodide in 10 ml NMP. The reaction mixture was stirred for two hours at rt. EtOAc was added and the organic layer was washed 5 times with brine and twice with water. The organic layer was reduced to 50% of volume, where upon 5-Benzo[1,3]dioxol-5-ylmethylene-2-methylsulfanyl-thiazol-4-one started to crystallize. Crystals were filtered off and washed with cold EtOAc.

Yield = 1.5 g (71%). HPLC: 3.44 min. LC-MS: M/Z ESI: 1.68 min, 280.19 (M+1). NMR: ^1H NMR (DMSO- d_6) δ 7.76 (s, 1H), 7.09-7.26 (m, 3H), 6.14 (s, 2H), 2.82 (s, 3H).

The following intermediates were synthesized according to the preparation of intermediate 9 using suitable starting materials.


Intermediate 10: Preparation of 2-Methylsulfanyl-5-quinoxalin-6-ylmethylene-thiazol-4-one

one



HPLC: 3.32 min. LC-MS: M/Z ESI: 1.20 min, 286.10 (M-1). NMR: ^1H NMR (DMSO- d_6) δ 9.01 (s, 2H), 8.31 (s, 1H), 8.21 (d, $J=8.7\text{Hz}$, 1H), 8.05 (d, $J=8.7\text{Hz}$, 1H), 7.90 (s, 1H), 2.85 (s, 3H).


Intermediate 11: Preparation of 2-Methylsulfanyl-5-quinolin-6-ylmethylene-thiazol-4-one



HPLC: 2.00 min. LC-MS: M/Z ESI: 1.48 min, 287.10 (M+1). NMR: ^1H NMR (DMSO- d_6) δ 8.97 (m, 1H), 8.50 (s, 1H), 8.12 (d, $J=8.7\text{Hz}$, 1H), 7.97-8.01 (m, 2H), 7.64 (dd, $J=4.1\text{Hz}$, 8.3Hz, 1H), 2.86 (s, 3H).

5 The following examples were synthesized:


Example 1: Preparation of N-(5-Benzo[1,3]dioxol-5-ylmethylene)-4-oxo-thiazolidin-2-ylidene)-2-chloro-benzenesulfonamide



5-Benzo[1,3]dioxol-5-ylmethylene-2-imino-thiazolidin-4-one (100mg, 0.4mmol) were dissolved in 3 ml NMP, followed by diisopropylethylamine (250ul) and 2-chlorobenzene-sulfonylchloride. After 8 hours the reaction was complete. Ethylacetate was added and the organic layer was washed with brine and dried over MgSO_4 . The crude was purified on Parallax Flex. Yield = 33.0mg (17%). HPLC: 3.92 min. LC-MS: M/Z ESI: 1.49 min, 421.06 (M-1). NMR: ^1H NMR (DMSO- d_6) δ 13.1 (b s, 1H), 8.12 (d, $J=9\text{Hz}$, 1H), 7.69-7.72 (m, 3H), 7.61 (m, 1H), 7.16- 7.21 (m, 2H), 7.13 (d, $J=9\text{Hz}$, 1H), 6.15 (s, 2H).


The following compounds were synthesized according to the synthesis of example 1 using suitable aldehydes such as e.g. piperonal, 6-quinolinecarboxaldehyde, 6-quinoxalinecarboxaldehyde, 2,2-difluoro-1,3-benzodioxole-5-carboxaldehyde:

Example 2: Preparation of ethanesulfonic acid (5-benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-amide




Yield = 50.0mg (37%). HPLC: 3.04 min. LC-MS: M/Z ESI: 1.49 min, 339.16 (M-1).

5 Example 3: Preparation of N-(5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-3-chloro-benzenesulfonamide




Yield = 30.0mg (16%). HPLC: 4.15 min. LC-MS: M/Z ESI: 1.54 min, 421.15 (M-1).

10 Example 4: Preparation of 5-Chloro-1,3-dimethyl-1H-pyrazole-4-sulfonic acid (5-benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-amide



Yield = 41.0mg (23%). HPLC: 3.58 min. LC-MS: M/Z ESI: 1.35 min, 439.05 (M-1).


Example 5: Preparation of 3-(5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidenesulfamoyl)-thiophene-2-carboxylic acid methyl ester



Yield = 46.0mg (25%). HPLC: 3.66 min. LC-MS: M/Z ESI: 1.38 min, 439.05 (M-1).


NMR: ^1H NMR (DMSO- d_6) δ 13.0 (b s, 1H), 8.0 (d, J = 3Hz, 1H), 7.74 (s, 1H), 7.59 (d, J = 3Hz, 1H), 7.13- 7.24 (m, 3H), 6.15 (s, 2H), 3.82 (s, 3H).

5 Example 6: Preparation of 6-Chloro-pyridine-3-sulfonic acid (5-benzofuran-2-ylmethylene-4-oxo-thiazolidin-2-ylidene)-amide




Yield = 21.0mg (13%). HPLC: 3.78 min. LC-MS: M/Z ESI: 1.46 min, 422.05 (M-1).

10 Example 7: Preparation of Quinoline-8-sulfonic acid (5-benzofuran-2-ylmethylene-4-oxo-thiazolidin-2-ylidene)-amide




Yield = 27.0mg (15%). HPLC: 3.59 min. LC-MS: M/Z ESI: 1.38 min, 438.04 (M-1).

Example 8: Preparation of N-(5-Benzofuran-2-ylmethylene-4-oxo-thiazolidin-2-ylidene)-benzenesulfonamide




Yield = 80.0mg (51%). HPLC: 3.82 min. LC-MS: M/Z ESI: 1.42 min, 387.11 (M-1).
NMR: ^1H NMR (DMSO- d_6) δ 13.5 (b s, 1H), 7.89 (m, 2H), 7.60-7.65 (m, 4H), 7.15-7.20 (m, 3H), 6.15 (s, 2H).

- 5 Example 9: Preparation of N-(5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-4-methyl-benzenesulfonamide




Yield = 90.0mg (52%). HPLC: 4.00 min. LC-MS: M/Z ESI: 1.51 min, 401.11 (M-1).

- 10 Example 10: Preparation of N-(5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-methanesulfonamide



Yield = 16.0mg (13%). HPLC: 2.85 min. LC-MS: M/Z ESI: 1.17 min, 325.06 (M-1).


- Example 11: Preparation of N-(5-(2,2-Difluoro-benzo[1,3]dioxol-5-ylmethylene)-4-oxo-thiazolidin-2-ylidene)-benzenesulfonamide



Yield = 91.0mg (54%). HPLC: 4.33 min. LC-MS: M/Z ESI: 1.66 min, 423.24 (M-1).

NMR: ^1H NMR (DMSO- d_6) δ 13.2 (b s, 1H), 7.28-7.93 (m, 9H).


5 Example 12: Preparation of N-[5-(2,2-Difluoro-benzof[1,3]dioxol-5-ylmethylene)-4-oxo-thiazolidin-2-ylidene]-4-methyl-benzenesulfonamide



Yield = 90.0mg (53%). HPLC: 4.52 min. LC-MS: M/Z ESI: 1.65 min, 437.23 (M-1).

NMR: ^1H NMR (DMSO- d_6) δ 12.6 (b s, 1H), 7.30-7.96 (m, 8H), 2.15 (s, 3H).


10 Example 13: Preparation of N-[5-(2,2-Difluoro-benzof[1,3]dioxol-5-ylmethylene)-4-oxo-thiazolidin-2-ylidene]-methanesulfonamide



Yield = 18.0mg (12%). HPLC: 3.55 min. LC-MS: M/Z ESI: 1.39 min, 361.16 (M-1).

NMR: ^1H NMR (DMSO- d_6) δ 12.9 (b s, 1H), 7.43-7.96 (m, 4H), 3.15 (s, 3H).


Example 14: Preparation of Biphenyl-2-sulfonic acid (5-benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-amide



Yield = 28.0mg (15%). HPLC: 4.27 min. LC-MS: M/Z ESI: 1.70 min, 463.12 (M-1).

- 5 NMR: ^1H NMR (DMSO- d_6) δ 12.6 (b s, 1H), 7.97 (d, J = 6Hz, 1H), 7.44-7.55 (m, 3H), 6.91-7.16 (m, 9H), 5.98 (s, 2H).


Example 15: Preparation of Pyridine-3-sulfonic acid (5-benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-amide



- 10 Yield = 56.0mg (36%). HPLC: 3.22 min. LC-MS: M/Z ESI: 1.30 min, 388.12 (M-1).


NMR: ^1H NMR (DMSO- d_6) δ 9.08 (s, 1H), 8.90 (d, 1H), 8.31 (d, J = 8.2Hz, 1H), 7.62-7.81 (m, 2H), 7.10-7.30 (m, 3H), 6.16 (s, 2H).

Example 16: Preparation of 3-(4-Oxo-5-quinolin-6-ylmethylene-thiazolidin-2-ylidenesulfamoyl)-thiophene-2-carboxylic acid methyl ester




Yield = 137.0mg (74%). HPLC: 2.46 min. LC-MS: M/Z ESI: 1.33 min, 458.12 (M-1).
 NMR: ^1H NMR (DMSO- d_6) δ 9.06 (s, 1H), 8.55 (d, $J=8.2\text{Hz}$, 1H), 8.30 (s, 1H), 8.20 (d, $J=8.2\text{Hz}$, 1H), 8.03 (m, 3H), 7.62-7.70 (m, 2H), 3.80 (s, 3H).

5 Example 17: Preparation of 2-Chloro-N-(4-oxo-5-quinolin-6-ylmethylene-thiazolidin-2-ylidene)-benzenesulfonamide



Yield = 60.0mg (35%). HPLC: 2.74 min. LC-MS: M/Z ESI: 1.40 min, 428.09 (M-1).
 NMR: ^1H NMR (DMSO- d_6) δ 9.00 (s, 1H), 8.55 (d, $J=8.2\text{Hz}$, 1H), 8.32 (s, 1H), 8.00-8.20 (m, 4H), 8.03 (m, 3H), 7.50-7.72 (m, 4H).


Example 18: Preparation of 3-(5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidenesulfamoyl)-thiophene-2-carboxylic acid



35 mg (0.08mmol) of 3-(5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidenesulfamoyl)-thiophene-2-carboxylic acid methyl ester (example 5) were dissolved in THF/water. 6 mg of LiOH.H₂O were added, and the reaction was followed by TLC. After complete saponification the reaction medium was acidified to pH 3.5, where upon the desired compound precipitated. Washing and drying afforded 25mg (70%) of 3-(5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidenesulfamoyl)-thiophene-2-carboxylic acid.

HPLC: 3.20 min. LC-MS: M/Z ESI: 1.05 min, 393.09 (M-1). NMR: ¹H NMR (DMSO-d₆) δ 13.2 (b s, 2H), 7.94 (d, J = 3Hz, 1H), 7.71 (s, 1H), 7.55 (d, J = 3Hz, 1H), 7.13- 7.19 (m, 3H), 6.14 (s, 2H).

Example 19: Preparation of 5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene-cyanamide




200mg (0.72mmol) of 5-Benzo[1,3]dioxol-5-ylmethylene-2-methylsulfanyl-thiazol-4-one were dissolved in NMP to which was added a solution of potassium tert.butoxide in hexane (1.1eq.). The colour changed to orange. To this was added as a solid cyanamide (1.2eq.). The reaction was heated at 80°C under Ar. for 3h. HPLC indicated complete transformation. 150 ml EtOAc were added and washed with 0.1N HCl twice. The organic layer was then washed extensively with brine. The solvent was dried and evaporated to dryness leading to a yellowish solid. The crude was purified on Parallel Flex system affording a yellow solid, which was dissolved in THF followed by 1equivalent of 1N KOH. 20ml of water were added and the frozen solution was lyophilized yielding 107mg (51%) of 5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene-cyanamide as potassium salt.

HPLC: 2.97 min. LC-MS: M/Z ESI: 1.30 min, 272.06 (M-1). NMR: ^1H NMR (DMSO- d_6) (potassium salt). δ 7.36 (s, 1H), 6.98-7.07 (m, 4H), 6.08 (s, 2H).


The following compounds were synthesized according to the preparation of example 19 using suitable starting materials:

- 5 Example 20: Preparation of 5-Benzo[1,3]dioxol-5-ylmethylene-thiazolidine-2,4-dione 2-(O-methyl-oxime).




HPLC: 3.34 min. LC-MS: M/Z ESI: 1.61 min, 277.19 (M-1). NMR: ^1H NMR (DMSO- d_6) (parent cpd). δ 12.1 (s, 1H), 7.55 (s, 1H), 7.05-7.14 (m, 3H), 6.12 (s, 2H), 3.80 (s, 3H).

- 10 Example 21: Preparation of 4-Oxo-5-quinoxalin-6-ylmethylene-thiazolidin-2-ylidene-cyanamide



HPLC: 2.51 min. LC-MS: M/Z ESI: 1.07 min, 280.09 (M-1). NMR: ^1H NMR (DMSO- d_6) (parent cpd). δ 12.80 (b s, 1H), 9.00 (s, 2H), 8.05-8.32 (m, 4H).


- 15 Example 22: 5-Benzo[1,3]dioxol-5-ylmethylene-2-benzylimino-thiazolidin-4-one



5-Benzo[1,3]dioxol-5-ylmethylene-2-thioxo-thiazolidin-4-one (100mg, 0.37mmol) were dissolved in EtOH/water, followed by 60mg of Na_2CO_3 . The reaction was stirred for 30min and benzylamine (122ul, 3 eq.) was added. The reaction mixture was refluxed for 2h. and the solvents were evaporated. The crude was purified by Parallel Flex chromatography. Yield: 31 mg (23%). HPLC: 3.66 min. LC-MS: M/Z ESI: 1.58 min, 339.13 (M+1). ^1H NMR (DMSO-d_6) δ 10.0 (s, 1H), 7.54 (s, 1H), 7.07-7.37 (m, 8H), 6.10 (s, 2H), 4.72 (s, 2H).


The following compounds were prepared according to the synthesis of example 24 using suitable starting materials:

Example 23: Preparation of 2-Benzylimino-5-quinolin-6-ylmethylene-thiazolidin-4-one.




Yield: 40 mg (27%). HPLC: 2.32 min. LC-MS: M/Z ESI: 1.52 min, 346.13 (M+1).

Example 24: Preparation of 2-Propylimino-5-quinolin-6-ylmethylene-thiazolidin-4-one.




Yield: 100 mg (80%). HPLC: 2.45 min. LC-MS: M/Z ESI: 1.32 min, 298.03 (M+1).

Example 25: 5-Benzo[1,3]dioxol-5-ylmethylene-2-propylimino-thiazolidin-4-one

Yield: 22 mg (14%). HPLC: 2.11 min. LC-MS: M/Z ESI: 1.46 min, 291.03 (M+1).

5 **Example 26: Preparation of 5-(4-Dimethylamino-quinazolin-6-ylmethylene)-2-methylimino-thiazolidin-4-one**



Yield: 42 mg (22%). HPLC: 1.48 min. LC-MS: M/Z ESI: 1.17 min, 314.05 (M+1).

The HPLC, NMR and MS data provided in the examples described below were obtained as followed: HPLC: column Waters Symmetry C8 50 x 4.6 mm, Conditions: MeCN/H₂O, 5 to 100% (8 min), max plot 230-400 nm; Mass spectra: PE-SCIEX API 150 EX (APCI and ESI), LC/MS spectra: Waters ZMD (ES); ¹H-NMR: Bruker DPX-300MHz.

The purifications were obtained as followed: Parallel Flex Biotage, Preparative HPLC Waters Prep LC 4000 System equipped with columns Prep Nova-Pak[®]HR C186 μm 60Å, 40x30mm (up to 100mg) or 40x300 mm (up to 1g). All the purifications were performed with a gradient of MeCN/H₂O 0.09% TFA.

Example 27 : Preparation of a pharmaceutical formulation

The following formulation examples illustrate representative pharmaceutical compositions according to the present invention being not restricted thereto.

Formulation 1 – Tablets

A compound of formula (I) is admixed as a dry powder with a dry gelatin binder in an approximate 1:2 weight ratio. A minor amount of magnesium stearate is added as a lubricant. The mixture is formed into 240-270 mg tablets (80-90 mg) of active 2-imino-azolinone compound per tablet) in a tablet press.

Formulation 2 – Capsules

A compound of formula (I) is admixed as a dry powder with a starch diluent in an approximate 1:1 weight ratio. The mixture is filled into 250 mg capsules (125 mg of active 2-imino-azolinone compound per capsule).

Formulation 3 – Liquid

A compound of formula (I) (1250 mg), sucrose (1.75 g) and xanthan gum (4 mg) are blended, passed through a No. 10 mesh U.S. sieve, and then mixed with a previously prepared solution of microcrystalline cellulose and sodium carboxymethyl cellulose (11:89, 50 mg) in water. Sodium benzoate (10 mg), flavor, and color are diluted with water and added with stirring. Sufficient water is then added to produce a total volume of 5 mL.

Formulation 4 – Tablets

A compound of formula (I) is admixed as a dry powder with a dry gelatin binder in an approximate 1:2 weight ratio. A minor amount of magnesium stearate is added as a lubricant. The mixture is formed into 450-900 mg tablets (150-300 mg of active 2-imino-azolinone compound) in a tablet press.

Formulation 5 – Injection

A compound of formula (I) is dissolved in a buffered sterile saline injectable aqueous medium to a concentration of approximately 5 mg/mL.

Example 28 : Biological assays

The compounds of the present invention may be subjected to the following assays:

5 a) High Throughput PI3K lipid kinase assay (binding assay):

The assay combines the scintillation proximity assay technology (SPA, Amersham) with the capacity of neomycin (a polycationic antibiotic) to bind phospholipids with high affinity and specificity. The Scintillation Proximity Assay is based on the properties of weakly emitting isotopes (such as ^3H , ^{125}I , ^{32}P). Coating SPA beads with neomycin allows
10 the detection of phosphorylated lipid substrates after incubation with recombinant PI3K and radioactive ATP in the same well, by capturing the radioactive phospholipids to the SPA beads through their specific binding to neomycin.

To a 384 wells MTP containing 5 μl of the test compound of formula (I) (solubilized in 6% DMSO; to yield a concentration of 100, 30, 10, 3, 1, 0.3, 0.1, 0.03, 0.01, 0.001 μM of the
15 test compound), the following assay components are added. 1) 5 μl (58 ng) of Human recombinant GST-PI3K γ (in Hepes 40 mM, pH 7.4, DTT 1 mM and ethylenglycol 5%) 2) 10 μl of lipid micelles and 3) 10 μl of Kinase buffer ($[\text{P}^{32}]\gamma\text{-ATP}$ 45 μM /60nCi, MgCl_2 30mM, DTT 1mM, β -Glycerophosphate 1mM, Na_3VO_4 100 μM , Na Cholate 0.3 %, in Hepes 40 mM, pH 7.4). After incubation at room temperature for 180 minutes, with gentle
20 agitation, the reaction is stopped by addition of 60 μl of a solution containing 100 μg of neomycin-coated PVT SPA beads in PBS containing ATP 10mM and EDTA 5mM. The assay is further incubated at room temperature for 60 minutes with gentle agitation to allow binding of phospholipids to neomycin-SPA beads. After precipitation of the neomycin-coated PVT SPA beads for 5 minutes at 1500 x g, radioactive *PutIns(3)P* is quantified by
25 scintillation counting in a Wallac MicroBeta TM plate counter.

The values indicated in respect of PI3K γ refer to the IC₅₀ (μ M), i.e. the amount necessary to achieve 50% inhibition of said target. Said values show a considerable potency of the 2-imino-azolinone-vinyl fused-benzene compounds with regard to PI3K γ .

The tested compounds according to formula (I) display an inhibition (IC₅₀) with regard to PI3K γ of less than 10 μ M, more preferred equal or less than 1 μ M.

Examples of inhibitory activities for test compounds 16, 20, 21, 22 & 26 are set out in Table 1.

<i>Example No</i>	<i>PI3Kγ, IC₅₀ (μ M)</i>
20	< 1.0
21	< 1.0
26	< 1.0
16	<1.0
22	< 1.5

Table 1: IC₅₀ values of 2-imino-azolinone-vinyl fused-benzene derivatives against PI3K γ .

b) Cell based ELISA to monitor PI3K inhibition:

Measurement of Akt/PKB phosphorylation in macrophages after stimulation with C5a:
 Raw 264: Raw 264-7 macrophages (cultured in DMEM-F12 medium containing 10% Fetal Calf serum and antibiotics) are plated at 20'000 cells/well in a 96 MTP 24 h before cell stimulation. Previous to the stimulation with 50 nM of Complement 5a during 5 minutes, Cells are serum starved for 2h, and pretreated with inhibitors for 20 minutes. After stimulation cells are fixed in 4% formaldehyde for 20 minutes and washed 3 times in PBS containing 1% Triton X-100 (PBS/Triton). Endogenous peroxidase is blocked by a 20 minutes incubation in 0.6% H₂O₂ and 0.1% Sodium Azide in PBS/Triton and washed 3 times in PBS/Triton. Cells are then blocked by 60 minutes incubation with 10% fetal calf

serum in PBS/Triton. Next, phosphorylated Akt/PKB is detected by an overnight incubation at 4°C with first antibody (anti phospho Serine 473 Akt IHC, Cell Signaling) diluted 800-fold in PBS/Triton, containing 5% bovine serum albumin (BSA). After 3 washes in PBS/Triton, cells are incubated for 60 minutes with a peroxidase conjugated
5 goat-anti-rabbit antibody (1/400 dilution in PBS/Triton, containing 5% BSA), washed 3 times in PBS/Triton, and 2 times in PBS and further incubated in 100 µl of substrate reagent solution (R&D) for 20 minutes. The reaction is stopped by addition of 50 µl of 1 M SO_4H_2 and absorbance is read at 450 nm.

The values indicated reflect the percentage of inhibition of AKT phosphorylation as
10 compared to basal level. Said values show a clear effect of the 2-imino-azolinone-vinyl fused-benzene compounds on the activation of AKT phosphorylation in macrophages.


For example the compound of example 8, when used at 10µM, completely inhibits (100 % inhibition) C5a-mediated AKT phosphorylation. The compound of example 10, when
15 used at 10 µM, inhibits 80% of the C5a-mediated AKT-phosphorylation.

Reference List

1. Vanhaesebroeck B et al., *Trends Biochem Sci.* **22**(7) p.267-72 (1997);
2. Leslie N.R et al *Chem Rev.* **101**(8) p.2365-80 (2001);
- 5 3. Katso R. et al. *Annu Rev Cell Dev Biol.* **17** p.615-75 (2001)
4. Toker a. et al *Cell Mol Life Sci.* **59**(5) p.761-79 (2002);
5. Vanhaesebroeck B. *Exp Cell Res.* **25**(1) p.239-54 (1999)
6. Stein R.C *Mol Med Today* **6**(9) p.347-57 (2000);
7. Wyman NP et al. *Immunol Today* **21**(6) p.260-4 (2000);
- 10 8. Hirsch et al *Science* **287**(5455) p.1049-53 (2000);
9. Hirsch et al. *FASEB J* **15**(11) p.2019-21 (2001);
10. Gerard C. et al. *Nat Immunol.* **2**(2) p.108-15 (2001);
11. Panayotou et al., *Trends Cell Biol.* **2** p.358-60 (1992);
12. Parker et al., *Current Biology*, **5** p.577-99 (1995),
- 15 13. Yao et al., *Science*, **267** p.2003-05 (1995);
14. Pages et al., *Nature*, **369** p.327-29 (1994);
15. Rudd, *Immunity* **4** p.527-34 (1996).
16. Fraser et al., *Science*, **251** p.313-16 (1991);
17. *J. Biol. Chem.* **273**(5) p.2505-8 (1998);
- 20 18. Lawlor MA et al (*J.Cell. Sci.* **114**(Pt 16) p.2903-10 (2001);
19. Laffargue M. et al *Immunity* **16**(3) p.441-51 (2002);
20. Stephens L. et al. *Curr. Opinion Cell Biol.* **14**(2) p.203-13 (2002);
21. Fruman et al., *An. Rev. Biochem.*, **67** p.481-507 (1998);
22. Thelen et al., *Proc. Natl. Acad. Sci. USA*, **91** p.4960-64 (1994);
- 25 23. Brummond et.al., *J.O.C.*, **64**, 1723-1726 (1999);
24. John M. Janusz et al, *J.Med.Chem.* 1998; Vol 41, No. 18.

Claims

1. An imino-azolinone-vinyl fused-benzene derivative according to formula (I),



S

X is S, O or -NR³:

Y is S or O:

10

15


20

alkynyl aryl or -heteroaryl, carboxy, cyano, hydroxy, C₁-C₆-alkoxy, nitro, acylamino, ureido, sulfonylamino, sulfanyl, or sulfonyl;

G is a C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, heteroaryl, C₁-C₆-alkyl aryl, C₁-C₆-alkyl heteroaryl, C₂-C₆-alkenyl-aryl or -heteroaryl, C₂-C₆-alkynyl aryl or -heteroaryl, C₁-C₆-alkoxy, cyano, C₁-C₆-acyl, or a sulfonyl moiety.

R³ is selected from the group comprising or consisting of H or C₁-C₆-alkyl.


with the proviso that the following 4 compounds are excluded :



2. The imino-azolinone-vinyl fused-benzene derivative according to claim 1, wherein A is selected from the group consisting of 2H-(benzo-1, 3-dioxolanyl), 2H, 3H-benzo-1,4-dioxanyl, 2,3-dihydrobenzofuranyl, anthraquinonyl, 2,2-difluorobenzo-1,3-dioxolenyl, 1,3-dihydrobenzofuranyl, benzofuranyl, 4-methyl-2H-benzo-1,4-oxazin-3-onyl, pyridinyl, 4-methyl-2H, 3H-benzo-1,4-oxazinyl.

3. The imino-azolinone-vinyl fused-benzene derivative according to claim 2, wherein A is a dioxolenyl or a pyridinyl moiety.

4. The imino-azolinone-vinyl fused-benzene derivative according to any of the preceding claims wherein R^1 and/or R^2 are H.
5. The imino-azolinone-vinyl fused-benzene derivative according to any of the preceding claims wherein G is a sulfonyl moiety of the formula $-SO_2-R^4$, whereby R^4 is selected from the group comprising or consisting of H, C_1-C_6 -alkyl, C_2-C_6 -alkenyl, C_2-C_6 -alkynyl, C_1-C_6 -alkyl carboxy, C_1-C_6 -alkyl acyl, C_1-C_6 -alkyl alkoxycarbonyl, C_1-C_6 -alkyl aminocarbonyl, C_1-C_6 -alkyl acyloxy, C_1-C_6 -alkyl acylamino, C_1-C_6 -alkyl ureido, C_1-C_6 -alkyl carbamate, C_1-C_6 -alkyl amino, C_1-C_6 -alkyl alkoxy, C_1-C_6 -alkyl sulfanyl, C_1-C_6 -alkyl sulfinyl, C_1-C_6 -alkyl sulfonyl, C_1-C_6 -alkyl sulfonylaminoaryl, aryl, heteroaryl, C_3-C_6 -cycloalkyl or heterocycloalkyl, C_1-C_6 -alkyl aryl, C_1-C_6 -alkyl heteroaryl, C_2-C_6 -alkenyl-aryl or -heteroaryl, C_2-C_6 -alkynyl aryl or -heteroaryl, carboxy, hydroxy, C_1-C_6 -alkoxy, acylamino, sulfonylamino.
6. The imino-azolinone-vinyl fused-benzene derivative according to claim 5, wherein R^4 is aryl, heteroaryl or C_1-C_3 alkyl.
7. The imino-azolinone-vinyl fused-benzene derivative according to any of the preceding claims wherein X is S, Y is O, R^1 and R^2 are H, A is a dioxolenyl or pyridinyl moiety.
8. An imino-azolinone-vinyl fused-benzene derivative according to formula (I)



wherein A is an 5-8 membered heterocyclic group or an carbocyclic group which may be fused with an aryl, an heteroaryl, an cycloalkyl or an heterocycloalkyl;

X is S, O or $-NR^3$;

Y is S or O;

5 R^1 is selected from the group comprising or consisting of H, CN, carboxy, acyl, C_1-C_6 -alkoxy, halogen, hydroxy, acyloxy, C_1-C_6 -alkyl carboxy, C_1-C_6 -alkyl acyloxy, C_1-C_6 -alkyl alkoxy, alkoxy carbonyl, C_1-C_6 -alkyl alkoxy carbonyl, aminocarbonyl, C_1-C_6 -alkyl aminocarbonyl, acylamino, C_1-C_6 -alkyl acylamino, ureido, C_1-C_6 -alkyl ureido, amino, C_1-C_6 -alkyl amino, ammonium, sulfonyloxy, C_1-C_6 -alkyl sulfonyloxy,
10 sulfonyl, C_1-C_6 -alkyl sulfonyl, sulfinyl, C_1-C_6 -alkyl sulfinyl, sulfanyl, C_1-C_6 -alkyl sulfanyl, sulfonylamino, C_1-C_6 -alkyl sulfonylamino or carbamate;

R^2 is selected from the group comprising or consisting of H, halogen, acyl, amino, C_1-C_6 -alkyl, C_2-C_6 -alkenyl, C_2-C_6 -alkynyl, C_1-C_6 -alkyl carboxy, C_1-C_6 -alkyl acyl, C_1-C_6 -alkyl alkoxy carbonyl, C_1-C_6 -alkyl aminocarbonyl, C_1-C_6 -alkyl acyloxy, C_1-C_6 -alkyl acylamino, C_1-C_6 -alkyl ureido, C_1-C_6 -alkyl carbamate, C_1-C_6 -alkyl amino, C_1-C_6 -alkyl alkoxy, C_1-C_6 -alkyl sulfanyl, C_1-C_6 -alkyl sulfinyl, C_1-C_6 -alkyl sulfonyl, C_1-C_6 -alkyl sulfonylaminoaryl, aryl, heteroaryl, C_3-C_8 -cycloalkyl or heterocycloalkyl, C_1-C_6 -alkyl aryl, C_1-C_6 -alkyl heteroaryl, C_2-C_6 -alkenyl-aryl or -heteroaryl, C_2-C_6 -alkynyl aryl or -heteroaryl, carboxy, cyano, hydroxy, C_1-C_6 -alkoxy, nitro, acylamino,
20 ureido, sulfonylamino, sulfanyl, or sulfonyl;

G is a C_1-C_6 -alkyl, C_2-C_6 -alkenyl, C_2-C_6 -alkynyl, heteroaryl, C_1-C_6 -alkyl aryl, C_1-C_6 -alkyl heteroaryl, C_2-C_6 -alkenyl-aryl or -heteroaryl, C_2-C_6 -alkynyl aryl or -heteroaryl, C_1-C_6 -alkoxy, cyano, C_1-C_6 -acyl, or a sulfonyl moiety.

R^3 is selected from the group comprising or consisting of H or C_1-C_6 -alkyl,

for use as a medicament.

9. The imino-azolinone-vinyl fused-benzene derivative according to any of the preceding claims, selected from the group consisting of :

N-(5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-2-chloro-benzenesulfonamide

Ethanesulfonic acid (5-benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-amide

N-(5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-3-chloro-benzenesulfonamide

5-Chloro-1,3-dimethyl-1H-pyrazole-4-sulfonic acid (5-benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-amide

3-(5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidenesulfamoyl)-thiophene-2-carboxylic acid methyl ester

Preparation of 6-Chloro-pyridine-3-sulfonic acid (5-benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-amide

Quinoline-8-sulfonic acid (5-benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-amide

N-(5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-benzenesulfonamide

N-(5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-4-methyl-benzenesulfonamide

N-(5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-methanesulfonamide

N-[5-(2,2-Difluoro-benzo[1,3]dioxol-5-ylmethylene)-4-oxo-thiazolidin-2-ylidene]-benzenesulfonamide

N-[5-(2,2-Difluoro-benzo[1,3]dioxol-5-ylmethylene)-4-oxo-thiazolidin-2-ylidene]-4-methyl-benzenesulfonamide

N-[5-(2,2-Difluoro-benzo[1,3]dioxol-5-ylmethylene)-4-oxo-thiazolidin-2-ylidene]-methanesulfonamide

Biphenyl-2-sulfonic acid (5-benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-amide

Pyridine-3-sulfonic acid (5-benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene)-amide

3-(4-Oxo-5-quinolin-6-ylmethylene-thiazolidin-2-ylidenesulfamoyl)-thiophene-2-carboxylic acid methyl ester

2-Chloro-N-(4-oxo-5-quinolin-6-ylmethylene-thiazolidin-2-ylidene)-benzenesulfonamide

3-(5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidenesulfamoyl)-thiophene-2-carboxylic acid

5-Benzo[1,3]dioxol-5-ylmethylene-4-oxo-thiazolidin-2-ylidene-cyanamide

5-Benzo[1,3]dioxol-5-ylmethylene-thiazolidine-2,4-dione 2-(O-methyl-oxime)

Preparation of 4-Oxo-5-quinoxalin-6-ylmethylene-thiazolidin-2-ylidene-cyanamide

5-Benzo[1,3]dioxol-5-ylmethylene-2-benzylimino-thiazolidin-4-one


2-Benzylimino-5-quinolin-6-ylmethylene-thiazolidin-4-one

2-Propylimino-5-quinolin-6-ylmethylene-thiazolidin-4-one

5-Benzo[1,3]dioxol-5-ylmethylene-2-propylimino-thiazolidin-4-one

5-(4-Dimethylamino-quinazolin-6-ylmethylene)-2-methylamino-thiazol-4-one

10. Use of an imino-azolinone-vinyl fused-benzene derivative according to formula (I)



wherein A is an 5-8 membered heterocyclic group or an carbocyclic group which may be fused with an aryl, an heteroaryl, an cycloalkyl or an heterocycloalkyl;

X is S, O or $-NR^3$;

Y is S or O;

R¹ is selected from the group comprising or consisting of H, CN, carboxy, acyl, C₁-C₆-alkoxy, halogen, hydroxy, acyloxy, C₁-C₆-alkyl carboxy, C₁-C₆-alkyl acyloxy, C₁-C₆-alkyl alkoxy, alkoxycarbonyl, C₁-C₆-alkyl alkoxycarbonyl, aminocarbonyl, C₁-C₆-alkyl aminocarbonyl, acylamino, C₁-C₆-alkyl acylamino, ureido, C₁-C₆-alkyl ureido, amino, C₁-C₆-alkyl amino, ammonium, sulfonyloxy, C₁-C₆-alkyl sulfonyloxy, sulfonyl, C₁-C₆-alkyl sulfonyl, sulfinyl, C₁-C₆-alkyl sulfinyl, sulfanyl, C₁-C₆-alkyl sulfanyl, sulfonylamino, C₁-C₆-alkyl sulfonylamino or carbamate;

R² is selected from the group comprising or consisting of H, halogen, acyl, amino, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₁-C₆-alkyl carboxy, C₁-C₆-alkyl acyl, C₁-C₆-alkyl alkoxy, C₁-C₆-alkyl alkoxycarbonyl, C₁-C₆-alkyl aminocarbonyl, C₁-C₆-alkyl acyloxy, C₁-C₆-alkyl acylamino, C₁-C₆-alkyl ureido, C₁-C₆-alkyl carbamate, C₁-C₆-alkyl amino, C₁-C₆-alkyl alkoxy, C₁-C₆-alkyl sulfanyl, C₁-C₆-alkyl sulfinyl, C₁-C₆-alkyl sulfonyl, C₁-C₆-alkyl sulfonylaminoaryl, aryl, heteroaryl, C₃-C₈-cycloalkyl or heterocycloalkyl, C₁-C₆-alkyl aryl, C₁-C₆-alkyl heteroaryl, C₂-C₆-alkenyl-aryl or -heteroaryl, C₂-C₆-alkynyl aryl or -heteroaryl, carboxy, cyano, hydroxy, C₁-C₆-alkoxy, nitro, acylamino, ureido, sulfonylamino, sulfanyl, or sulfonyl;

G is a C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, heteroaryl, C₁-C₆-alkyl aryl, C₁-C₆-alkyl heteroaryl, C₂-C₆-alkenyl-aryl or -heteroaryl, C₂-C₆-alkynyl aryl or -heteroaryl, C₁-C₆-alkoxy, cyano, C₁-C₆-acyl, or a sulfonyl moiety.

R³ is selected from the group comprising or consisting of H or C₁-C₆-alkyl,

5 for the preparation of a medicament for the prophylaxis and/or treatment of autoimmune disorders and/or inflammatory diseases, cardiovascular diseases, neurodegenerative diseases, bacterial or viral infections, kidney diseases, platelet aggregation, cancer, transplantation, graft rejection or lung injuries.

11. Use according to claim 10, wherein said diseases are selected in the group including
10 multiple sclerosis, psoriasis, rheumatoid arthritis, multiple sclerosis, systemic lupus erythematosus, inflammatory bowel disease, lung inflammation, thrombosis or brain infection/inflammation such as meningitis or encephalitis.

12. Use according to claim 10, wherein said diseases are selected in the group including
15 Alzheimer's disease, Huntington's disease, CNS trauma, stroke or ischemic conditions.

13. Use according to claim 10, wherein said diseases are selected in the group including atherosclerosis, heart hypertrophy, cardiac myocyte dysfunction, elevated blood pressure or vasoconstriction.

14. Use according to claim 10, wherein said diseases are selected in the group including
20 chronic obstructive pulmonary disease, anaphylactic shock fibrosis, psoriasis, allergic diseases, asthma, stroke or ischemic conditions, ischemia-reperfusion, platelets aggregation/activation, skeletal muscle atrophy/hypertrophy, leukocyte recruitment in cancer tissue, angiogenesis, invasion metastasis, in particular melanoma, Kaposi's sarcoma, acute and chronic bacterial and viral infections, sepsis, graft rejection,

glomerulo sclerosis, glomerulo nephritis, progressive renal fibrosis, endothelial and epithelial injuries in the lung or in general lung airways inflammation.


15. Use according to any of claims 10 to 14 for the modulation, in particular for the inhibition, of the PI3 kinase activity.
- 5 16. Use according to claim 15, wherein said PI3 kinase is a PI3 kinase γ .
17. A pharmaceutical composition containing at least one thiazolidinone-vinyl fused-benzene derivative according to any of claims 1 to 9 and a pharmaceutically acceptable carrier, diluent or excipient thereof.
18. A method of preparing a 2-imino-azolinone-vinyl fused-benzene derivatives of formula (I) according to any of claims 1 to 9 comprising the following step:
- 10



wherein A, R¹, R², G, X and Y are as above defined and L is a leaving group.

Abstract of the invention:

The present invention is related to 2-imino-azolinone-vinyl fused-benzene derivatives of formula (I) in particular for the treatment and/or prophylaxis of autoimmune disorders and/or inflammatory diseases, cardiovascular diseases, neurodegenerative diseases,
 5 bacterial or viral infections, kidney diseases, platelet aggregation, cancer, transplantation, graft rejection or lung injuries.



wherein A is an 5-8 membered heterocyclic group or an carbocyclic group which may be fused with an aryl, an heteroaryl, an cycloalkyl or an heterocycloalkyl;
 10

X is S, O or -NR³;

Y is S or O;

G is a C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, heteroaryl, C₁-C₆-alkoxy, cyano, C₁-C₆-acyl, or a sulfonyl moiety.